

APC anti-human CD83

Catalog # / Size: 305311 / 25 tests
305312 / 100 tests

Clone: HB15e


Isotype: Mouse IgG1, κ

Reactivity: Human, **Cross-Reactivity:** Chimpanzee, Baboon, Cynomolgus, Rhesus, Pigtailed Macaque

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Monocytes-derived dendritic cells
(induced with GM-CSF+IL-4+TNF- α)
stained with HB15e APC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁴.

Application References:

- Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press New York.
- Zhou L, *et al.* 1995. *J. Immunol.* 154:3821.
- Cao W, *et al.* 2005. *Biochem. J.* 385:85.
- Lore K, *et al.* 2002. *AIDS* 16:683. (IHC)
- Cho H, *et al.* 2007. *Physiol Genomics* doi:10.1152/physiolgenomics.00051.2006
- Joo H, *et al.* 2012. *J Exp Med.* PubMed.

Description: CD83 is a 43 kD single chain type I glycoprotein also known as HB15. A member of the immunoglobulin superfamily, CD83 is expressed on a subset of dendritic cells, Langerhans cells, and weakly on activated lymphocytes. Although CD83 is thought to play a role in antigen presentation and/or lymphocyte activation, the precise function of this protein is unknown. CD83 is considered to be a useful marker for mature dendritic cells.

Antigen References:

- Kozlow E, *et al.* 1993. *Blood* 81:454.
- Zhou L, *et al.* 1992. *J. Immunol.* 149:735.
- Zhou L, *et al.* 1995. *Blood* 86:3295.

Related Products:Product

APC anti-human CD1a	HI149
APC anti-human HLA-DR	L243
APC anti-human CD14	M5E2
APC Mouse IgG1, κ Isotype Ctrl	MOPC-21
Cell Staining Buffer	
RBC Lysis Buffer (10X)	
Human TruStain FcX™ (Fc Receptor Blocking Solution)	

Clone

HI149
L243
M5E2
MOPC-21

Application

FC
FC
FC
FC, ICFC
FC, ICC, ICFC
FC, ICFC
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.