

## **Product Data Sheet**

## Alexa Fluor® 488 anti-human CD83

Catalog # / Size: 305314 / 100 tests

Clone: HB15e

**Isotype:** Mouse IgG1,  $\kappa$ 

Reactivity: Human, Cross-Reactivity: Chimpanzee, Baboon, Cynomolgus, Rhesus,

Pigtailed Macaque

Preparation: The antibody was purified by affinity chromatography, and conjugated with


Alexa Fluor® 488 under optimal conditions. The solution is free of


unconjugated Alexa Fluor® 488.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from





Human monocytes-derived mature dendritic cells stained with HB15e Alexa Fluor® 488

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is 5  $\mu$ l per million cells or 5  $\mu$ l per 100  $\mu$ l of whole blood. It is recommended that the reagent be titrated for optimal

performance for each application.

\* Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at

\*\* Alexa Fluor® is a registered trademark of Molecular Probes, Inc. Alexa Fluor® dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and

issued patents.

frozen tissue sections<sup>4</sup>

1. Knapp W, et al. 1989. Leucocyte Typing IV. Oxford University Press New York. Application References:

2. Zhou L, et al. 1995. J. Immunol. 154:3821. 3. Cao W, et al. 2005. Biochem. J. 385:85. 4. Lore K, et al. 2002. AIDS 16:683. (IHC)

5. Cho H, et al. 2007. Physiol Genomics doi:10.1152/physiolgenomics.00051.2006

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed

**Description:** CD83 is a 43 kD single chain type I glycoprotein also known as HB15. A member of the immunoglobulin superfamily,

CD83 is expressed on a subset of dendritic cells, Langerhans cells, and weakly on activated lymphocytes. Although CD83 is thought to play a role in antigen presentation and/or lymphocyte activation, the precise function of this protein

is unknown. CD83 is considered to be a useful marker for mature dendritic cells.

Antigen References: 1. Kozlow E, et al. 1993. Blood 81:454.

2. Zhou L, et al. 1992. J. Immunol. 149:735.

3. Zhou L, et al. 1995. Blood 86:3295.

**Application Related Products: Product** Clone Cell Staining Buffer

RBC Lysis Buffer (10X)

FC, ICC, ICFC FC, ICFC Alexa Fluor® 488 Mouse IgG1, κ Isotype Ctrl (FC) MOPC-21 FC, IF Human TruStain FcX™ (Fc Receptor Blocking Solution) FC, ICC, ICFC



