

# Product Data Sheet

## Purified anti-human CD74

**Catalog # / Size:** 326802 / 100 µg

**Clone:** LN2

**Isotype:** Mouse IgG1, κ

**Immunogen:** SU-DHL-4 cells


**Reactivity:** Human, **Cross-Reactivity:** Baboon, Rhesus

**Preparation:** The antibody was purified by affinity chromatography.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C.



Human peripheral blood lymphocytes stained with purified LN2, followed by anti-mouse IgG FITC

## Applications:

**Applications:** FC - *Quality tested*  
IHC, IP - *Reported in the literature*

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 2.0 µg per 10<sup>6</sup> cells in 100 µl volume or 100 µl whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** Additional reported applications (for the relevant formats) include: immunohistochemical staining<sup>1,2</sup> of frozen sections and formalin-fixed paraffin-embedded sections<sup>1,2</sup>, and immunoprecipitation<sup>1</sup>.

**Application References:** 1. Epstein AL, *et al.* 1984. *J. Immunol.* 133:1028.  
2. Marder RJ, *et al.* 1985. *Lab. Invest.* 52:497.

**Description:** CD74 is a type II transmembrane glycoprotein also known as MHC class II associated invariant chain, invariant chain, Ii, MHC class II chaperone, and MIF receptor. CD74 exists in four isoforms with molecular masses of 33, 35, 41, and 43 kD, depending on genetic splicing. CD74 is primarily expressed on antigen presenting cells, including B cells, monocytes/macrophages, dendritic cells, and Langerhans cells. It is also expressed by activated T cells and activated endothelial and epithelial cells as well as carcinomas of lung, renal, gastric and thymic origin. The primary function of CD74 is intracellular sorting of MHC class II molecules and regulation of exogenous peptide loading onto MHC class II. It is also involved in the modulation of B cell differentiation and positive selection of CD4<sup>+</sup> T cells. It has been reported that CD74 binds MIF (macrophage migration inhibitory factor) and signals through CD44 to regulate innate and adaptive immunity. It is also reported that H. pylori infection occurs through urease B binding of CD74 on gastric epithelial cells, inducing gastric epithelial cell apoptosis, NF-κB activation, and IL-8 production.

**Antigen References:** 1. Moldenhauer G, *et al.* 1999. *Immunology* 96:473.  
2. Shi X, *et al.* 2006. *Immunity* 25:595.  
3. Beswick EJ, *et al.* 2006. *Infect. Immun.* 74:1148.  
4. Zola H, *et al.* 2007. *Leukocyte and Stromal Cell Molecules: The CD Markers* Wiley-Liss A John Wiley & Sons Inc, Publication

### Related Products:

**Product**  
Purified Mouse IgG1, κ Isotype Ctrl  
APC Goat anti-mouse IgG (minimal x-reactivity)  
PE Goat anti-mouse IgG (minimal x-reactivity)  
Cell Staining Buffer

**Clone**  
MOPC-21  
Poly4053  
Poly4053

**Application**  
FC, ICFC, ICC, IF, IHC, IP, WB  
FC  
FC  
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biollegend.com/ordering#license](http://www.biollegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.