

PE anti-human CD79a (Ig α)

Catalog # / Size: 333503 / 25 tests
333504 / 100 tests

Clone: HM47

Isotype: Mouse IgG1, κ


Workshop Number: V cB017

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes surface stained with CD19 (HIB19) FITC and intracellular stained with HM47 PE

Applications:

Applications: ICFC - *Quality tested*

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application References: 1. Mason DY, *et al.* 1991. *J. Immunol.* 147:2474
2. Bhargava P, *et al.* 2007. *Am. J. Clin. Pathol.* 128:306

Description: CD79a is a 47kD type I integral membrane protein, also known as mb-1 or Iga. It is a member of the Ig superfamily and disulphide-associated with CD79b (B29). The interaction of CD79a/CD79b heterodimer with B cell surface Ig forms B cell antigen complex. CD79a is expressed in B cells from early pre-B to plasma cell stage. It has been shown that CD79a is also weakly expressed in some precursors of T- and myeloid cells. CD79 mediates the transport of IgM to B cell surface and transduces signals initiated by BCR aggregation

Antigen References: 1. Zola Heddy, *et al.* Eds. 2007. *Leukocyte and Stromal Cell Molecules: The CD markers.* WILEY-LISS
2. Astsaturov IA, *et al.* 1996. *Leukemia* 10:769
3. Mason DY, *et al.* 1995 *Blood* 86:1453
4. Hashimoto M, *et al.* 2002. *J. Pathol.* 197:341

Related Products:

Product
PE Mouse IgG1, κ Isotype Ctrl
Permeabilization Wash Buffer (10X)
Cell Staining Buffer
RBC Lysis Buffer (10X)
Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone
MOPC-21

Application
FC, ICFC
ICC, ICFC, IHC
FC, ICC, ICFC
FC, ICFC
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.