

## **Product Data Sheet**

## Alexa Fluor® 647 anti-human CD82

performance for each application.

633nm / 635nm.


issued patents.


Catalog # / Size:	342108 / 100 tests		
Clone:	ASL-24		
Isotype:	Mouse IgG1, κ		
Reactivity:	Human		
Preparation:	The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 647 under optimal conditions. The solution is free of unconjugated Alexa Fluor® 647.		
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).		
Storage:	The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. <b>Do not freeze.</b>		
Applications:			
Applications:	FC - Quality tested		
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the		

suggested use of this reagent is 5  $\mu$ l per million cells or 5  $\mu$ l per 100  $\mu$ l of whole blood. It is recommended that the reagent be titrated for optimal

\* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at

\*\* Alexa Fluor® is a registered trademark of Molecular Probes, Inc. Alexa Fluor® dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and





FC, ICC, ICFC

**Description:** CD82 is a 45-90 kD type III tetraspan membrane protein which is encoded by the KAI1 gene. A member of the 4-span transmembrane protein superfamily (TM4SF) CD82 forms a complex with CD37, CD53, CD81, ECM and MHC molecules. CD82 is expressed on monocytes, granulocytes, lymphocytes, epithelial cells, endothelial cells, and fibroblasts and plays a role in signal transduction and adhesion. It has been suggested CD82 functions as a tumor suppressor as loss of expression has been found to promote tumor metastasis.

Antigen References: 1. Miranti CK. 2009. Cell. Signal. 21:196   2. Abe M, et al. 2008. 266:163 2. Abe M, et al. 2004. Cancer Res. 64:4235   3. Lee JH et al. 2004. Cancer Res. 64:4235 4. Lagaudriere-Gesbert C, et al. 1997. J. Immunol. 158:	:2790		
Related Products: Product Alexa Fluor® 647 Mouse IgG1, κ Isotype Ctrl (FC) Cell Staining Buffer PBC Lycis Buffer (10X)	Clone MOPC-21	Application FC, IF FC, ICC, ICFC FC, ICFC	

RBC Lysis Buffer (10X) Human TruStain FcX™ (Fc Receptor Blocking Solution)



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

(

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.