

Product Data Sheet

APC anti-human CD73 (Ecto-5'-nucleotidase)

Catalog # / Size: 344005 / 25 tests
344006 / 100 tests

Clone: AD2

Isotype: Mouse IgG1, κ


Workshop Number: V B-CD73.3

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with AD2 APC and CD19 PE (top) or mouse IgG1, κ APC isotype control and CD19 PE (bottom).

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application References: 1. Nakamura T, *et al.* 1993. *J. Immunol.* 151:6933.

Description: CD73 is a 70 kD glycoposphatidylinositol (GPI)-linked 5'-nucleotidase also known as ecto-5'-nucleotidase. It converts adenosine monophosphate (AMP) to adenosine. CD73 is expressed on subsets of T and B cells, mesenchymal stem cells, follicular dendritic cells, endothelial cells, and epithelial cells. It has been reported that CD73 costimulates T cell activation, and mediates adhesion of lymphocytes to follicular dendritic cells and endothelial cells.


Antigen References: 1. Zola H, *et al.* 2007. *Leukocyte and stromal Cell Molecules:the CD Markers.* A John Wiley & Sons Inc, Publication.
2. Airas L and Jalkanen S, *et al.* 1996. *Blood* 88:1755.
3. Gutensohn W, *et al.* 1995. *Cell Immunol.* 161:213.
4. Airas L, *et al.* 1995. *J. Exp. Med.* 182:1603.

Related Products:

Product
APC Mouse IgG1, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)
Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone
MOPC-21

Application
FC, ICFC
FC, ICC, ICFC
FC, ICFC
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.