

PE anti-human CD94

Catalog # / Size: 305506 / 100 tests

Clone: DX22

Isotype: Mouse IgG1, κ


Immunogen: NK cell clone

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with DX22 PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation⁴, inhibition of NK cell-mediated lysis⁵, and immunohistochemical staining of acetone-fixed frozen tissue sections.

Application References:

- Mizuki M, *et al.* 2000. *Sarcoidosis Vasc. Diffuse Lung Dis.* 17:54.
- Phillip J, *et al.* 1996. *Immunity* 5:163.
- Lazetic S, *et al.* 1996. *J. Immunol.* 157:4741.
- Lanier LL, *et al.* 1998. *Immunity* 8:693.
- Wooden SL, *et al.* 2005. *J. Immunol.* 175:1383.

Description: CD94 is a 43 kD type II transmembrane glycoprotein also known as KP43. CD94 belongs to the C-type lectin superfamily and is present as a covalently linked heterodimer with NKG2 on the cell surface. CD94 is expressed by NK cells, a subset of $\gamma\delta$ T cells, and NKT cells. The CD94/NKG2A complex serves as an inhibitory receptor specific for HLA-class I molecules.

Antigen References:

- Lopez-Botet M, *et al.* 1997. *Immunol. Rev.* 155:165.
- Moretta A, *et al.* 1997. *Immunol. Rev.* 155:105.

Related Products:	Product	Clone	Application
	PE anti-human CD16	3G8	FC
	PE Mouse IgG1, κ Isotype Ctrl	MOPC-21	FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	PE anti-human CD56 (NCAM)	MEM-188	FC
	Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.