

PE anti-human CD87

Catalog # / Size: 336905 / 25 tests
336906 / 100 tests

Clone: VIM5

Isotype: Mouse IgG1, κ

Workshop Number: VI MR13


Immunogen: THP-1 (human myeloid cell line)

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood granulocytes stained with VIM5 PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: Additional reported applications include: VIM5 mAb binds to uPA binding-domain

Application References:

- Schlossmant SF, *et al.* eds. 1995 *Leukocyte Type V*. Oxford University Press Inc, New York
- Gadd SJ, *et al.* 1990. *Clin. Exp. Immunol* 80:252
- Sillaber C, *et al.* 1997. *J. Biol. Chem.* 272:7824

Description: CD87, known as urokinase plasminogen activator-receptor (uPAR), is a 36-68 kD single-chain glycoposphotidylinositol (GPI)-anchored glycoprotein. The extracellular region of CD87 consists of three homologous domains, D1, D2, and D3. Membrane-bound and soluble (D2D3) forms of uPAR have been reported. The membrane-bound CD87 is expressed on granulocytes, monocytes/macrophages, dendritic cells, fibroblasts, endothelial cells and keratinocytes. CD87 plays a role in cell adhesion and chemotaxis through binding to β 1, β 2 and β 3 integrins. CD87-bound uPA converts plasminogen to plasmin.

Antigen References:

- Beaufort N, *et al.* 2004. *J. Immunol.* 172:540
- Tarui T, *et al.* 2001. *J. Bio. Chem.* 276:3983
- Zola H, *et al.* 2007. *Leukocyte and Stromal Cell Molecules: The CD Markers*. A John Wiley & Sons Inc, Publication

Related Products:	Product	Clone	Application
	PE Mouse IgG1, κ Isotype Ctrl (FC)	MOPC-21	FC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.