

## PE anti-human CD99

**Catalog # / Size:** 318008 / 100 tests

**Clone:** HCD99


**Isotype:** Mouse IgG2a,  $\kappa$

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Storage:** The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with HCD99 PE

## Applications:

**Applications:** FC - *Quality tested*

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20  $\mu$ l to 5  $\mu$ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100  $\mu$ l staining volume or per 100  $\mu$ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at [www.biolegend.com/testsize](http://www.biolegend.com/testsize) regarding the test size change.

**Application Notes:** Additional reported (for the relevant formats) applications include: immunohistochemical staining of acetone-fixed frozen sections and formalin-fixed paraffin-embedded tissues.

**Application References:** 1. Tadimitsu K, *et al.* eds. 1997. Leukocyte Typing VI Garland Publishing Inc. London.

**Description:** The HCD99 monoclonal antibody recognizes human CD99 also known as CD99 antigen, E2 antigen, MIC2, and T-cell surface glycoprotein E2. CD99 is a type I, single chain transmembrane protein devoid of N-linked glycosylation sites that is encoded by the pseudoautosomal gene MIC2. CD99 has an apparent molecular weight 32 kD and is widely expressed on a variety of tissues. CD99 is highly expressed on thymocytes, T cells, T cell leukemias and lymphomas and is absent on fetal B cells, some B cell lines, eosinophils, granulocytes and the NK-cell line YT. CD99 is involved in spontaneous rosette formation with erythrocytes and may also be involved in other T-cell and hematopoietic cell adhesion pathways. CD99 has been reported to activate a caspase-independent death pathway in T cells under some conditions. CD99 interacts with a number of proteins including ferritin heavy chain 1, karyopherin beta 1, TRIP13, cyclophilin A, annexin II, and ubiquitin-conjugating enzyme E2H. The HCD99 antibody has been reported to be useful for flow cytometric detection of human CD99 and immunohistochemistry (acetone-fixed frozen tissues and formalin-fixed paraffin-embedded tissues).

**Antigen References:** 1. Gelin C, *et al.* 1989. *EMBO J.* 8:3253.  
2. Goodfellow PJ, *et al.* 1986. *Science* 234:740.  
3. Pettersen RD, *et al.* 2001. *J. Immunol.* 166:4931.

### Related Products:Product

Cell Staining Buffer  
PE Mouse IgG2a,  $\kappa$  Isotype Ctrl (FC)  
Human TruStain FcX™ (Fc Receptor Blocking Solution)

### Clone

MOPC-173

### Application

FC, ICC, ICFC  
FC  
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.