


# Product Data Sheet

## Purified anti-human CD105

**Catalog # / Size:** 323202 / 100 µg  
**Clone:** 43A3  
**Isotype:** Mouse IgG1, κ  
**Immunogen:** L-cells transfected with human CD105  
**Reactivity:** Human  
**Preparation:** The antibody was purified by affinity chromatography.  
**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.  
**Concentration:** 0.5 mg/ml  
**Storage:** The antibody solution should be stored undiluted at 4°C.



Human monocytic cell line THP-1 stained with purified 43A3, followed by anti-mouse IgG FITC

## Applications:

**Applications:** FC - Quality tested

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤2.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application References:** 1. Bühring HJ, *et al.* 1991. *Leukemia* 5:841.  
 2. Vogel W, *et al.* 2002. *Haematologica* 88:126.

**Description:** CD105 is also known as Endoglin. It is a type I integral membrane homodimer protein with subunits of 90 kD found on vascular endothelial cells and syncytiotrophoblasts of placenta. CD105 is weakly expressed on stromal fibroblasts. It is also expressed on activated monocytes and tissue macrophages. Expression of CD105 is increased on activated endothelium in tissues undergoing angiogenesis, such as in tumors, or in cases of wound healing or dermal inflammation. CD105 is a component of the TGF-β receptor system in human umbilical vein endothelial cells and binds TGF-β1 and β3 with high affinity but does not bind to TGF-β2.

**Antigen References:** 1. Mason D, *et al.* Eds. 2002. *Leucocyte Typing VII*. Oxford University Press. New York.  
 2. Pierelli L, *et al.* 2001. *Leuk. Lymphoma* 42:1195.

### Related Products:

**Product**  
 Purified Mouse IgG1, κ Isotype Ctrl  
 APC Goat anti-mouse IgG (minimal x-reactivity)  
 FITC Goat anti-mouse IgG (minimal x-reactivity)  
 PE Goat anti-mouse IgG (minimal x-reactivity)  
 Cell Staining Buffer

**Clone**  
 MOPC-21  
 Poly4053  
 Poly4053  
 Poly4053

**Application**  
 FC, ICFC, ICC, IF, IHC, IP, WB  
 FC  
 FC  
 FC  
 FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biollegend.com/ordering#license](http://www.biollegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.