

APC anti-human CD105

Catalog # / Size: 323207 / 25 tests
323208 / 100 tests

Clone: 43A3

Isotype: Mouse IgG1, κ


Immunogen: L-cells transfected with human CD105

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human monocytic cell line THP-1
stained with 43A3 APC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application References:

1. Bühring HJ, *et al.* 1991. *Leukemia* 5:841.
2. Vogel W, *et al.* 2002. *Haematologica* 88:126.
3. Laviola L, *et al.* 2012. *Endocrinology*. 153:5770. PubMed.

Description: CD105 is also known as Endoglin. It is a type I integral membrane homodimer protein with subunits of 90 kD found on vascular endothelial cells and syncytiotrophoblasts of placenta. CD105 is weakly expressed on stromal fibroblasts. It is also expressed on activated monocytes and tissue macrophages. Expression of CD105 is increased on activated endothelium in tissues undergoing angiogenesis, such as in tumors, or in cases of wound healing or dermal inflammation. CD105 is a component of the TGF- β receptor system in human umbilical vein endothelial cells and binds TGF- β 1 and β 3 with high affinity but does not bind to TGF- β 2.

Antigen References:

1. Mason D, *et al.* Eds. 2002. Leucocyte Typing VII. Oxford University Press. New York.
2. Pierelli L, *et al.* 2001. *Leuk. Lymphoma* 42:1195.

Related Products:	Product	Clone	Application
	Cell Staining Buffer		FC, ICC, ICFC
	APC Mouse IgG1, κ Isotype Ctrl (FC)	MOPC-21	FC
	Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.