

Applications	Reactivity	Sensitivity	MW (kDa)	Isotype
W IP IHC-P IHC-F IF-IC F	H (M) (R)	Endogenous	185 kDa	Rabbit IgG

Applications Key: W=Western Blotting IP=Immunoprecipitation IHC-P=Immunohistochemistry (Paraffin) IHC-F=Immunohistochemistry (Frozen) IF-IC=Immunofluorescence (Immunocytochemistry) F=Flow Cytometry

Reactivity Key: H=Human M=Mouse R=Rat

Species cross-reactivity is determined by western blot. Species enclosed in parentheses are predicted to react based on 100% sequence homology.

Protocols

Flow Cytometry Protocol

A. Solutions and Reagents

NOTE: Prepare solutions with purified water.

- 1X Phosphate Buffered Saline (PBS):** Dissolve 8 g NaCl, 0.2 g KCl, 1.44 g Na₂HPO₄ and 0.24 g KH₂PO₄ in 800 ml dH₂O. Adjust the pH to 7.4 with HCl and the volume to 1 L. Store at room temperature.
- Formaldehyde** (methanol free).
- 100% Methanol**
- Incubation Buffer:** Dissolve 0.5 g bovine serum albumin (BSA) in 100 ml 1X PBS. Store at 4 °C.

B. Fixation

- Collect cells by centrifugation and aspirate supernatant.
- Resuspend cells briefly in 0.5–1 ml PBS. Add formaldehyde to a final concentration of 2–4% formaldehyde.
- Fix for 10 min at 37 °C.
- Chill tubes on ice for 1 min.
- For extracellular staining with antibodies that do not require permeabilization, proceed to Section D, Step 1 or store cells in PBS with 0.1% sodium azide at 4 °C; for intracellular staining, proceed to permeabilization (Section C, Step 1).

C. Permeabilization

- Permeabilize cells by adding ice-cold 100% methanol slowly to pre-chilled cells, while gently vortexing, to a final concentration of 90% methanol. Alternatively, to remove fix prior to permeabilization, pellet cells by centrifugation and resuspend in 90% methanol.
- Incubate 30 min on ice.
- Proceed with immunostaining (Section D, Step 1) or store cells at –20 °C in 90% methanol.

D. Immunostaining

NOTE: Account for isotype matched controls for monoclonal antibodies or species matched IgG for polyclonal antibodies. Count cells using a hemocytometer or alternative method.

- Aliquot 0.5–1x10⁶ cells into each assay tube (by volume).
- Add 2–3 ml Incubation Buffer to each tube and rinse by centrifugation. Repeat.
- Resuspend cells in 100 µl Incubation Buffer per assay tube.
- Block in Incubation Buffer for 10 min at room temperature.

5. Add the unconjugated, biotinylated, or fluorochrome-conjugated primary antibody at the appropriate dilution to the assay tubes (see individual antibody datasheet for the appropriate dilution).
6. Incubate for 1 hr at room temperature.
7. Rinse as before in Incubation Buffer by centrifugation.
8. If using a fluorochrome-conjugated primary antibody, resuspend cells in 0.5 ml PBS and analyze on flow cytometer; for unconjugated or biotinylated primary antibodies, proceed to immunostaining (Section D, Step 9).
9. Resuspend cells in fluorochrome-conjugated secondary antibody or fluorochrome-conjugated avidin, diluted in Incubation Buffer at the recommended dilution.
10. Incubate for 30 min at room temperature.
11. Rinse as before in Incubation Buffer by centrifugation.
12. Resuspend cells in 0.5 ml PBS and analyze on flow cytometer; alternatively, for DNA staining, proceed to optional DNA stain (Section E, Step 1).

E. Optional DNA Stain

1. Resuspend cells in 0.5 ml of DNA dye (e.g. Propidium Iodide (PI)/RNase Staining Solution [#4087](#)).
2. Incubate for at least 5 min at room temperature.
3. Analyze cells in DNA stain on flow cytometer.

Immunohistochemistry Protocol (Frozen)

A. Solutions and Reagents

1. Xylene
2. Ethanol (anhydrous denatured, histological grade 100% and 95%)
3. Hematoxylin (optional)
4. **Fixative: For optimal fixative, please refer to the product data sheet**
 - a. 10% Neutral buffered formalin
 - b. Acetone
 - c. Methanol
 - d. 16% formaldehyde
1. **3% formaldehyde:** To prepare, add 18.75 ml 16% formaldehyde to 81.25 ml 1X PBS.
5. **10X Tris Buffered Saline (TBS):** To Prepare 1 L add 24.2 g Trizma base ($C_4H_{11}NO_3$) and 80 g sodium chloride (NaCl) to 1 L dH₂O. Adjust pH to 7.6 with concentrated HCl.
6. **Wash buffer:** 1X Tris Buffered Saline (TBS) To prepare 1 L add 100 ml 10X TBS to 900 ml dH₂O.
7. **Methanol/Peroxidase:** To prepare, add 10 ml 30% H₂O₂ to 90 ml methanol. Store at -20 °C.
8. **Blocking Solution:** 1X TBS/0.3% Triton-X 100/5% normal goat serum ([#5425](#)). **To prepare:** add 500 µl goat serum and 30 µl Triton-X 100 to 9.5 ml 1X TBS.
9. **Biotinylated Secondary Antibody.**
10. **ABC Reagent:** (Vectastain ABC Kit, Vector Laboratories, Inc., Burlingame, CA). Prepare according to manufacturer's instructions 30 minutes before use.

11. **DAB Reagent or suitable substrate:** Prepare according to manufacturer's recommendations.

B. Sectioning

1. For tissue stored at -80 °C: remove from freezer and equilibrate at -20 °C for approximately 15 minutes before attempting to section. This may prevent cracking of the block when sectioning.
2. Section tissue at a range of 6-8 µm and place on positively charged slides.
3. Allow sections to air dry on bench for a few minutes before fixing (this helps sections adhere to slides).

C. Fixation

NOTE: Consult product data sheet to determine the optimal fixative.

1. After sections have dried on the slide, fix in optimal fixative as directed below.
 - a. **10% Neutral buffered formalin:** 10 minutes at room temperature. Proceed with staining procedure immediately.
 - b. **Cold acetone:** 10 minutes at -20 °C. Air dry. Proceed with staining procedure immediately.
 - c. **Methanol:** 10 minutes at -20 °C. Proceed with staining procedure immediately.
 - d. **3% Formaldehyde:** 15 minutes at room temperature. Proceed with staining procedure immediately.
 - e. **3% Formaldehyde/methanol:** 15 minutes at room temperature in 3% formaldehyde, followed by 5 minutes in methanol at -20 °C (**do not rinse in between**). Proceed with staining procedure immediately.

D. Staining

1. Wash sections in wash buffer twice for 5 minutes.
2. Incubate for 10 minutes at room temperature in 3% H₂O₂ diluted in methanol.
3. Wash sections in wash buffer twice for 5 minutes.
4. Block each section with blocking solution for one hour at room temperature.
5. Remove blocking solution and add 100-400 µl diluted primary antibody to each section. (Dilute antibody in blocking solution). Incubate overnight at 4 °C. **Refer to product datasheet to determine the recommended dilution.*
6. Remove antibody solution and wash sections three times with wash buffer for 5 minutes each.
7. Add 100-400 µl secondary antibody, diluted in blocking solution per manufacturer's recommendation, to each section. Incubate 30 minutes at room temperature.
8. If using ABC avidin/biotin method, make ABC reagent according to the manufacturer's instructions and incubate solution for 30 minutes at room temperature.
9. Remove secondary antibody solution and wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 µl ABC reagent to each section and incubate for 30 min. at room temperature.
11. Remove ABC reagent and wash sections three times in wash buffer for 5 minutes each.
12. Add 100-400 µl DAB or suitable substrate to each section and monitor staining closely.
13. As soon as the sections develop, immerse slides in dH₂O.
14. If desired, counterstain sections in Hematoxylin per manufacturer's instructions.
15. Wash sections in dH₂O two times for 5 minutes each.
16. **Dehydrate sections:**
 - a. Incubate sections in 95% ethanol two times for 10 seconds each.

- b. Repeat in 100% ethanol, incubating sections two times for 10 seconds each.
 - c. Repeat in xylene, incubating sections two times for 10 seconds each.
17. Mount coverslips.

Immunohistochemistry Protocol (Paraffin)

***IMPORTANT:** See product data sheet for the appropriate antibody diluent and antigen unmasking procedure. **IHC**

Protocol: Unmasking buffer/antibody diluent.

A. Solutions and Reagents

1. Xylene
2. Ethanol, anhydrous denatured, histological grade (100% and 95%)
3. Deionized water (dH₂O)
4. Hematoxylin (optional)
5. **Wash Buffer:**

1X TBS/0.1% Tween-20 (1X TBST): To prepare 1 L add 100 ml 10X TBS to 900 ml dH₂O. Add 1 ml Tween-20 and mix.

10X Tris Buffered Saline (TBS): To prepare 1 L add 24.2 g Trizma[®] base (C₄H₁₁NO₃) and 80 g sodium chloride (NaCl) to 1 L dH₂O. Adjust pH to 7.6 with concentrated HCl.
6. ***Antibody Diluent:**
 - a. **SignalStain[®] Antibody Diluent #8112**
 - b. **TBST/5% normal goat serum (#5425):** To 5 ml 1X TBST add 250 µl normal goat serum.
 - c. **PBST/5% normal goat serum (#5425):** To 5 ml 1X PBST add 250 µl normal goat serum.

1X PBS/0.1% Tween-20 (1X PBST): To prepare 1 L add 100 ml 10X PBS to 900 ml dH₂O. Add 1 ml Tween-20 and mix.

10X Phosphate Buffered Saline (PBS): To prepare 1 L add 80 g sodium chloride (NaCl), 2 g potassium chloride (KCl), 14.4 g sodium phosphate, dibasic (Na₂HPO₄) and 2.4 g potassium phosphate, monobasic (KH₂PO₄) to 1 L dH₂O. Adjust pH to 7.4.
7. ***Antigen Unmasking:**
 - a. **Citrate:** 10 mM Sodium Citrate Buffer: To prepare 1 L add 2.94 g sodium citrate trisodium salt dihydrate (C₆H₅Na₃O₇•2H₂O) to 1 L dH₂O. Adjust pH to 6.0.
 - b. **EDTA:** 1 mM EDTA: To prepare 1 L add 0.372 g EDTA (C₁₀H₁₄N₂O₈Na₂•2H₂O) to 1 L dH₂O. Adjust pH to 8.0.
 - c. **TE:** 10 mM Tris/1 mM EDTA, pH 9.0: To prepare 1L add 1.21 g Trizma[®] base (C₄H₁₁NO₃) and 0.372 g EDTA (C₁₀H₁₄N₂O₈Na₂•2H₂O) to 950 ml dH₂O. Adjust pH to 9.0, then adjust final volume to 1000 ml with dH₂O.
 - d. **Pepsin:** 1 mg/ml in Tris-HCl pH 2.0.
8. **3% Hydrogen Peroxide:** To prepare, add 10 ml 30% H₂O₂ to 90 ml dH₂O.
9. **Blocking Solution:** TBST/5% normal goat serum (#5425): to 5 ml 1X TBST add 250 µl normal goat serum.
10. Biotinylated secondary antibody.
11. **ABC Reagent:** (Vectastain ABC Kit, Vector Laboratories, Inc., Burlingame, CA) Prepare according to manufacturer's instructions 30 minutes before use.

12. **DAB Reagent or suitable substrate:** Prepare according to manufacturer's recommendations.

B. Deparaffinization/Rehydration

NOTE: Do not allow slides to dry at any time during this procedure.

1. **Deparaffinize/hydrate sections:**
 - a. Incubate sections in three washes of xylene for 5 minutes each.
 - b. Incubate sections in two washes of 100% ethanol for 10 minutes each.
 - c. Incubate sections in two washes of 95% ethanol for 10 minutes each.
2. Wash sections twice in dH₂O for 5 minutes each.

C. *Antigen Unmasking

NOTE: Consult product data sheet for specific recommendation for the unmasking solution.

1. **For Citrate:** Bring slides to a boil in 10 mM sodium citrate buffer pH 6.0 then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench top for 30 minutes.
2. **For EDTA:** Bring slides to a boil in 1 mM EDTA pH 8.0 followed by 15 minutes at a sub-boiling temperature. No cooling is necessary.
3. **For TE:** Bring slides to a boil in 10 mM TE/1 mM EDTA, pH 9.0 then maintain at a sub-boiling temperature for 18 minutes. Cool on the bench for 30 minutes.
4. **For Pepsin:** Digest for 10 minutes at 37 °C.

D. Staining

1. Wash sections in dH₂O three times for 5 minutes each.
2. Incubate sections in 3% hydrogen peroxide for 10 minutes.
3. Wash sections in dH₂O twice for 5 minutes each.

NOTE: Consult product data sheet for recommended antibody diluent.
4. Wash sections in wash buffer for 5 minutes.
5. Block each section with 100-400 µl blocking solution for 1 hour at room temperature.
6. Remove blocking solution and add 100-400 µl primary antibody diluted in recommended antibody diluent to each section. Incubate overnight at 4 °C.
7. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
8. Add 100-400 µl biotinylated secondary antibody, diluted in TBST per manufacturer's recommendation, to each section. Incubate 30 minutes at room temperature.
9. If using ABC avidin/biotin method, prepare ABC reagent according to the manufacturer's instructions and incubate solution for 30 minutes at room temperature.
10. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
11. Add 100-400 µl ABC reagent to each section and incubate for 30 minutes at room temperature.
12. Remove ABC reagent and wash sections three times in wash buffer for 5 minutes each.
13. Add 100-400 µl DAB or suitable substrate to each section and monitor staining closely.
14. As soon as the sections develop, immerse slides in dH₂O.

15. If desired, counterstain sections in hematoxylin per manufacturer's instructions.
16. Wash sections in dH₂O two times for 5 minutes each.
17. Dehydrate sections:
 - a. Incubate sections in 95% ethanol two times for 10 seconds each.
 - b. Repeat in 100% ethanol, incubating sections two times for 10 seconds each.
 - c. Repeat in xylene, incubating sections two times for 10 seconds each.
18. Mount coverslips.

Immunofluorescence Protocol for Methanol Fixation (IF Methanol-fixed)

IMPORTANT: Please refer to the **APPLICATIONS** section on the front page of the datasheet to determine if this product is validated and approved for use with this protocol.

A. Solutions and Reagents

NOTE: Prepare solutions with Milli-Q or equivalently purified water.

- **10X Phosphate Buffered Saline (PBS):** To prepare 1 L add 80 g sodium chloride (NaCl), 2 g potassium chloride (KCl), 14.4 g sodium phosphate, dibasic (Na₂HPO₄) and 2.4 g potassium phosphate, monobasic (KH₂PO₄) to 1 L dH₂O. Adjust pH to 8.0.
- **Methanol, 100%**
- **Blocking Buffer** (1X PBS / 5% normal goat serum ([#5425](#)) / 0.3% Triton X-100):
To prepare 25 ml, add 2.5 ml 10X PBS, 1.25 ml normal serum from the same species as the secondary antibody (eg. normal goat serum, normal donkey serum) and 21.25 ml dH₂O and mix well. While stirring, add 75 µl Triton X-100.
- **Antibody Dilution Buffer** (1X PBS / 1% BSA / 0.3% Triton X-100):
To prepare 40 ml, add 4 ml 10X PBS to 36 ml dH₂O, mix. Add 0.4 g BSA and mix well. While stirring, add 120 µl Triton X-100.
- **Fluorochrome-conjugated secondary antibody** ([recommended secondary antibodies](#))
NOTE: When using any primary or fluorochrome-conjugated secondary antibody for the first time, titrate the antibody to determine which dilution allows for the strongest specific signal with the least background for your sample.
- **Prolong[®] Gold Antifade Reagent** ([Invitrogen](#), Eugene, OR, cat# P36930)

B. Specimen Preparation - Cultured Cell Lines (IF-IC)

NOTE: Cells should be grown, treated, fixed and stained directly in multi-well plates, chamber slides or on coverslips.

1. Aspirate liquid, then cover cells to a depth of 2–3 mm with ice-cold 100% methanol.
2. Allow cells to fix for 15 minutes at -20C.
3. Aspirate fixative, rinse three times in PBS for 5 minutes each.
4. Proceed with Immunostaining (Section C).

C. Immunostaining

NOTE: All subsequent incubations should be carried out at room temperature unless otherwise noted in a humid light-tight box or covered dish/plate to prevent drying and fluorochrome fading.

1. Block specimen in Blocking Buffer for 60 minutes.

2. While blocking, prepare primary antibody by diluting as indicated on datasheet in Antibody Dilution Buffer.
 3. Aspirate blocking solution, apply diluted primary antibody.
 4. Incubate overnight at 4 °C.
 5. Rinse three times in PBS for 5 minutes each.
- NOTE:** If using primary antibodies directly conjugated with Alexa Fluor® fluorochromes, skip to step C8.
6. Incubate specimen in fluorochrome-conjugated [secondary antibody](#)* diluted in Antibody Dilution Buffer for 1–2 hours at room temperature in dark.
 7. Rinse in PBS as in step 5.
 8. Coverslip slides with Prolong® Gold Antifade Reagent.
 9. For best results, examine specimens immediately using appropriate excitation wavelength. For long-term storage, store slides flat at 4 °C protected from light.

Immunoprecipitation Protocol / (For Analysis By Western Immunoblotting)

For **shorter assay times** please try our [Immunoprecipitation Protocol Utilizing Magnetic Separation / \(For Analysis By Western Immunoblotting\)](#).

A. Solutions and Reagents

NOTE: Prepare solutions with Milli-Q or equivalently purified water.

1. 1X Phosphate Buffered Saline (PBS)
 2. **1X Cell Lysis Buffer:** ([#9803](#)) 20 mM Tris (pH 7.5), 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% Triton X-100, 2.5 mM Sodium pyrophosphate, 1 mM β-glycerophosphate, 1 mM Na₃VO₄, 1 μg/ml Leupeptin
- NOTE:** Add 1 mM PMSF immediately prior to use.
3. **Protein A or G Agarose Beads:** (Protein A [#9863](#)) Please prepare according to manufacturer's instructions. Use Protein A for rabbit IgG pull down and Protein G for mouse IgG pull down.
 4. **3X SDS Sample Buffer:** ([#7722](#)) 187.5 mM Tris-HCl (pH 6.8 at 25 °C), 6% w/v SDS, 30% glycerol, 150 mM DTT, 0.03% w/v bromophenol blue

B. Preparing Cell Lysates

1. Aspirate media. Treat cells by adding fresh media containing regulator for desired time.
2. To harvest cells under nondenaturing conditions, remove media and rinse cells once with ice-cold PBS.
3. Remove PBS and add 0.5 ml ice-cold 1X cell lysis buffer to each plate (10 cm) and incubate the plates on ice for 5 minutes.
4. Scrape cells off the plates and transfer to microcentrifuge tubes. Keep on ice.
5. Sonicate samples on ice three times for 5 seconds each.
6. Microcentrifuge for 10 minutes at 14,000 X g, 4 °C, and transfer the supernatant to a new tube. If necessary, lysate can be stored at –80 °C.

C. Immunoprecipitation

Optional: It may be necessary to perform a lysate pre-clearing step to reduce non-specific binding to the Protein A/G agarose beads (See section below).

1. Take 200 μ l cell lysate and add primary antibody. Incubate with gentle rocking overnight at 4 $^{\circ}$ C.
2. Add either protein A or G agarose beads (20 μ l of 50% bead slurry). Incubate with gentle rocking for 1–3 hours at 4 $^{\circ}$ C.
3. Microcentrifuge for 30 seconds at 4 $^{\circ}$ C. Wash pellet five times with 500 μ l of 1X cell lysis buffer. Keep on ice during washes.
4. Resuspend the pellet with 20 μ l 3X SDS sample buffer. Vortex, then microcentrifuge for 30 seconds.
5. Heat the sample to 95–100 $^{\circ}$ C for 2–5 minutes and microcentrifuge for 1 minute at 14,000 X g.
6. Load the sample (15–30 μ l) on SDS-PAGE gel (12–15%).
7. Analyze sample by Western blotting (see Western Immunoblotting Protocol: [Western BSA](#), [Western Milk](#)).

Cell Lysate Pre-Clearing (Optional)

1. Take 200 μ l cell lysate and add to either Protein A or G agarose beads (20 μ l of 50% bead slurry).
2. Incubate at 4 $^{\circ}$ C for 30 – 60 minutes.
3. Spin for 10 minutes at 4 $^{\circ}$ C. Transfer the supernatant to a fresh tube.
4. Proceed to step 1 of Immunoprecipitation.

NOTE: For proteins with molecular weights of 50 kDa, we recommend using [Mouse Anti-Rabbit IgG \(Light-Chain Specific\) \(L57A3\) mAb #3677](#) or [Mouse Anti-Rabbit IgG \(Conformation Specific\) \(L27A9\) mAb #3678](#) as a secondary antibody to minimize masking produced by denatured heavy chains. For proteins with molecular weights of 25 kDa, [Mouse Anti-Rabbit IgG \(Conformation Specific\) \(L27A9\) mAb #3678](#) is recommended.

Western Immunoblotting Protocol (Primary Ab Incubation In BSA)

For Western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4 $^{\circ}$ C with gentle shaking, overnight.

Products available from Cell Signaling Technology are linked by their respective catalog numbers.

A. Solutions and Reagents

NOTE: Prepare solutions with Milli-Q or equivalently purified water.

1. **1X Phosphate Buffered Saline (PBS).**
2. **1X SDS Sample Buffer:** ([#7722](#), [#7723](#)) 62.5 mM Tris-HCl (pH 6.8 at 25 $^{\circ}$ C), 2% w/v SDS, 10% glycerol, 50 mM DTT, 0.01% w/v bromophenol blue or phenol red.
3. **Transfer Buffer:** 25 mM Tris base, 0.2 M glycine, 20% methanol (pH 8.5).
4. **10X Tris Buffered Saline (TBS):** ([#9997](#)) To prepare 1 liter of 10X TBS: 24.2 g Tris base, 80 g NaCl; adjust pH to 7.6 with HCl (use at 1X).
5. **Nonfat Dry Milk:** ([#9999](#)) (weight to volume [w/v]).
6. **Blocking Buffer:** 1X TBS, 0.1% Tween-20 with 5% w/v nonfat dry milk; for 150 ml, add 15 ml 10X TBS to 135 ml water, mix. Add 7.5 g nonfat dry milk and mix well. While stirring, add 0.15 ml Tween-20 (100%).
7. **Wash Buffer:** 1X TBS, 0.1% Tween-20 (TBS/T).
8. **Bovine Serum Albumin (BSA):** ([#9998](#)).
9. **Primary Antibody Dilution Buffer:** 1X TBS, 0.1% Tween-20 with 5% BSA; for 20 ml, add 2 ml 10X TBS to 18 ml water, mix. Add 1.0 g BSA and mix well. While stirring, add 20 μ l Tween-20 (100%).

10. **Phototope[®]-HRP Western Blot Detection System:** ([#7071 anti-rabbit](#)) or ([#7072 anti-mouse](#)) Includes biotinylated protein ladder, secondary ([#7074 anti-rabbit](#)) or ([#7076 anti-mouse](#)) antibody conjugated to horseradish peroxidase (HRP), anti-biotin antibody conjugated to HRP, LumiGLO[®] chemiluminescent reagent and peroxide.
11. **Prestained Protein Marker, Broad Range (Premixed Format):** ([#7720](#)).
12. **Biotinylated Protein Ladder Detection Pack:** ([#7727](#)).
13. **Blotting Membrane:** This protocol has been optimized for nitrocellulose membranes, which CST recommends. PVDF membranes may also be used.

B. Protein Blotting

A general protocol for sample preparation is described below.

1. Treat cells by adding fresh media containing regulator for desired time.
2. Aspirate media from cultures; wash cells with 1X PBS; aspirate.
3. Lyse cells by adding 1X SDS sample buffer (100 µl per well of 6-well plate or 500 µl per plate of 10 cm diameter plate). Immediately scrape the cells off the plate and transfer the extract to a microcentrifuge tube. Keep on ice.
4. Sonicate for 10–15 seconds for complete cell lysis and to shear DNA (to reduce sample viscosity).
5. Heat a 20 µl sample to 95–100 °C for 5 minutes; cool on ice.
6. Microcentrifuge for 5 minutes.
7. Load 20 µl onto SDS-PAGE gel (10 cm x 10 cm). **NOTE:** CST recommends loading prestained molecular weight markers ([#7720](#), 10 µl/lane) to verify electrotransfer and biotinylated protein ladder ([#7727](#), 10 µl/lane) to determine molecular weights.
8. Electrotransfer to nitrocellulose or PVDF membrane.

C. Membrane Blocking and Antibody Incubations

NOTE: Volumes are for 10 cm x 10 cm (100 cm²) of membrane; for different sized membranes, adjust volumes accordingly.

1. (Optional) After transfer, wash nitrocellulose membrane with 25 ml TBS for 5 minutes at room temperature.
2. Incubate membrane in 25 ml of blocking buffer for 1 hour at room temperature.
3. Wash three times for 5 minutes each with 15 ml of TBS/T.
4. Incubate membrane and primary antibody (at the appropriate dilution) in 10 ml primary antibody dilution buffer with gentle agitation overnight at 4 °C.
5. Wash three times for 5 minutes each with 15 ml of TBS/T.

I. For Unconjugated Primary Antibodies

1. Incubate membrane with **appropriate** HRP-conjugated secondary antibody (1:2000) and HRP-conjugated anti-biotin antibody (1:1000) to detect biotinylated protein markers in 10 ml of blocking buffer with gentle agitation for 1 hour at room temperature.
2. Wash three times for 5 minutes each with 15 ml of TBS/T.

II. For HRP Conjugated Primary Antibodies

Skip to Detection of Proteins (Step D).

III. For Biotinylated Primary Antibodies

1. Incubate membrane with HRP-Streptavidin (at the appropriate dilution) in milk for one hour with gentle agitation at room temperature.
2. Wash three times for 5 minutes each with 15 ml of TBS/T.

D. Detection of Proteins

1. Incubate membrane with 10 ml LumiGLO[®] (0.5 ml 20X LumiGLO[®], 0.5 ml 20X Peroxide and 9.0 ml Milli-Q water) with gentle agitation for 1 minute at room temperature. **NOTE:** LumiGLO[®] substrate can be further diluted if signal response is too fast.
2. Drain membrane of excess developing solution (do not let dry), wrap in plastic wrap and expose to x-ray film. An initial 10-second exposure should indicate the proper exposure time. **NOTE:** Due to the kinetics of the detection reaction, signal is most intense immediately following LumiGLO[®] incubation and declines over the following 2 hours.


Specificity / Sensitivity

HER2/ErbB2 (29D8) Rabbit mAb detects endogenous levels of total ErbB2 protein. This antibody does not cross-react with related kinases.

Source / Purification


Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding tyrosines 1248 of human ErbB2 protein.

Western Blotting



Western blot analysis of cell extracts from various cell lines, using HER2/ErbB2 (29D8) Rabbit mAb.


IHC-P (paraffin)



Immunohistochemical analysis of paraffin-embedded human breast carcinoma, using HER2/ErbB2 (29D8)

Rabbit mAb.


IHC-P (paraffin)



Immunohistochemical analysis of paraffin-embedded human breast carcinoma using HER2/ErbB2 (29D8)

RmAb in the presence of control peptide (left) or HER2/ErbB2 Blocking Peptide #1059 (right).


IHC-P (paraffin)



Immunohistochemical analysis of paraffin-embedded SKBr3 (high HER2) (left), MDA-MB-453 (moderate


HER2) (middle) and MDA-MB-468 (low HER2) (right), using HER2/ErbB2 (29D8) Rabbit mAb.

IHC-F (frozen)




Immunohistochemical analysis of frozen SKOV-3 xenograft using HER2/ErbB2 (29D8) Rabbit mAb.

Flow Cytometry



Flow cytometric analysis of MCF7 cells using HER2/ErbB2 (29D8) Rabbit mAb (blue) compared to a nonspecific negative control antibody (red).

IF-IC



Confocal immunofluorescent analysis of MDA-MB-453 cells (left) and MDA-MB-231 cells (right), using HER2/ErbB2 (29D8) Rabbit mAb (green). Blue pseudocolor = DRAQ5® #4084 (fluorescent DNA dye).