


Anti-Mouse Siglec H eFluor® 660 (Alexa® 647 Replacement)

Catalog Number: 50-0333

Also known as: Sialic Acid-Binding Immunoglobulin-Like Lectin H RUO: For Research Use Only. Not for use in diagnostic procedures.



Description

Host/Isotype: Rat IgG2b

The monoclonal antibody eBio440c recognizes Siglec-H, a protein exclusively found on pDC (plasmacytoid dendritic cells) or type I IFN-producing cells (IPC) in the naïve mouse. Mouse IPC are typically PDCA+, CD11c+, CD11b-, B220+, and Ly-6C+, and are quick to respond to viruses. Siglec-H is a transmembrane protein of the Ig superfamily that like CD33 have been shown to bind sialic acid but lacks the characteristic cytoplasmic ITIM domain (immunoreceptor tyrosine based inhibitory motif). To overcome the lack of a cytoplasmic domain, Siglec-H associates with DAP12 thereby allowing for signal transduction.

Use By: Refer to vial Contains sodium azide

The eBio440c antibody has been shown to inhibit pDC function (inhbits IFNalpha secretion in response to CpG).

It has been observed that some mouse strains (such as SJL) have higher percentages of pDCs compared to C57BL/6.

Applications Reported

This eBio440c antibody has been reported for use in flow cytometric analysis.

Applications Tested

This eBio440c antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.25 μ g per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

eFluor® 660 is a replacement for Alexa Fluor® 647. eFluor® 660 emits at 659 nm and is excited with the red laser (633 nm). Please make sure that your instrument is capable of detecting this fluorochome.

References

Kreisel FH, Blasius A, Kreisel D, Colonna M, Cella M. Interferon-producing cells develop from murine CD31(high)/Ly6C(-) marrow progenitors. Cell Immunol. 2006 Aug;242(2):91-8.



Anti-Mouse Siglec H eFluor® 660 (Alexa® 647 Replacement)

Catalog Number: 50-0333

Also known as: Sialic Acid-Binding Immunoglobulin-Like Lectin H RUO: For Research Use Only. Not for use in diagnostic procedures.

Blasius AL, Cella M, Maldonado J, Takai T, Colonna M. Siglec-H is an IPC-specific receptor that modulates type I IFN secretion through DAP12. Blood. 2006 Mar 15;107(6):2474-6.(440c, FC, PubMed)

Blasius AL, Giurisato E, Cella M, Schreiber RD, Shaw AS, Colonna M. Bone marrow stromal cell antigen 2 is a specific marker of type I IFN-producing cells in the naive mouse, but a promiscuous cell surface antigen following IFN stimulation. J Immunol. 2006 Sep 1;177(5):3260-5.(440c, FC, PubMed)

Blasius A, Vermi W, Krug A, Facchetti F, Cella M, Colonna M. A cell-surface molecule selectively expressed on murine natural interferon alpha producing cells that blocks secretion of interfero nalpha. Blood. 2004;103:4201-4206 (440c, FC, IH/F, FA PubMed)

Related Products

11-0452 Anti-Human/Mouse CD45R (B220) FITC (RA3-6B2) 12-3172 Anti-Mouse CD317 (BST2, PDCA-1) PE (eBio927) 50-4031 Rat IgG2b Isotype Control eFluor® 660 (Alexa Fluor® 647 Replacement)