


Anti-Mouse CD127 eFluor® 660 (Alexa Fluor® 647 Replacement)

Catalog Number: 50-1271

Also known as: Interleukin-7 Receptor alpha, IL-7Ra

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of C57Bl/6 splenocytes with Anti-Mouse CD3e FITC (cat. 11-0031) and 0.5 ug of Rat IgG2a K Isotype Control eFluor® 660 (cat. 50-4321) (left) or 0.5 ug of Anti-Mouse CD127 eFluor® 660 (right). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse CD127 eFluor® 660 (Alexa Fluor® 647 Replacement)

Catalog Number: 50-1271

Clone: A7R34

Concentration: 0.2 mg/mL

Host/Isotype: Rat IgG2a, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light-sensitive material.

Batch Code: Refer to vial

Use By: Refer to vial

Contains sodium azide



Description

The A7R34 monoclonal antibody reacts with mouse CD127, the high affinity alpha subunit of the mouse IL-7 receptor. IL-7 receptor alpha chain is expressed by immature B cells in the bone marrow, double-negative (CD4-CD8-), single-positive (CD4+ and CD8+), but not double-positive (CD4+CD8+) thymocytes. In the periphery, mature T cells express CD127 at low level. A7R34 inhibits binding of IL-7 to its receptor and has been used in *in vivo* and *in vitro* studies to elucidate the role of IL-7 in T and B cell development and activation. Binding of A7R34 blocks the binding of SB/199, another antibody which recognizes mouse CD127.

Applications Reported

This A7R34 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This A7R34 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 1 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

eFluor® 660 is a replacement for Alexa Fluor® 647. eFluor® 660 emits at 659 nm and is excited with the red laser (633 nm). Please make sure that your instrument is capable of detecting this fluorochrome.

References

Sudo, T., S. Nishikawa, et al. 1993. Expression and function of the interleukin 7 receptor in murine lymphocytes. *Proc Natl Acad Sci U S A* 90(19): 9125-9.

Ohana M, Okazaki K, Oshima C, et al. 2001. A critical role for IL-7R signaling in the development of *Helicobacter felis*-induced gastritis in mice. *Gastroenterology*. Aug;121(2):329-36.

Okuno Y, Iwasaki H, Huettner CS, Radomska HS, Gonzalez DA, Tenen DG, Akashi K. 2002. Differential regulation of

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com • info@ebioscience.com

Anti-Mouse CD127 eFluor® 660 (Alexa Fluor® 647 Replacement)

Catalog Number: 50-1271

Also known as: Interleukin-7 Receptor alpha, IL-7Ra

RUO: For Research Use Only. Not for use in diagnostic procedures.

the human and murine CD34 genes in hematopoietic stem cells. Proc Natl Acad Sci U S A. Apr 30;99(9):6246-51.

Leithauser F, Meinhardt-Krajina T, et al. 2006. Foxp3-expressing CD103+ regulatory T cells accumulate in dendritic cell aggregates of the colonic mucosa in murine transfer colitis. Am J Pathol. 168(6):1898-909. (**A7R34**, IHC frozen, PubMed)

LTbetaR signaling induces cytokine expression and up-regulates lymphangiogenic factors in lymph node anlagen. Vondenhoff MF, Greuter M, Goverse G, Elewaut D, Dewint P, Ware CF, Hoorweg K, Kraal G, Mebius RE. J Immunol. 2009 May 1;182(9):5439-45. (**A7R34**, IF, PubMed)

Related Products

11-0031 Anti-Mouse CD3e FITC (145-2C11)

12-5773 Anti-Mouse/Rat Foxp3 PE (FJK-16s)

50-4321 Rat IgG2a K Isotype Control eFluor® 660 (eBR2a)