
Anti-Mouse CD8a eFluor[®] 650NC (for IHC/ICC)


Catalog Number: IH95-0081

Also known as: CD8 alpha, Ly-2, Ly-35, Ly-B, Lyt-2

RUO: For Research Use Only. Not for use in diagnostic procedures.

Product Information

Contents: Anti-Mouse CD8a eFluor[®] 650NC
(for IHC/ICC)

 **Catalog Number:** IH95-0081

Clone: 53-6.7

Host/Isotype: Rat IgG2a, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light-sensitive material.

Batch Code: Refer to vial

Use By: Refer to vial







Description

The 53-6.7 monoclonal antibody reacts with the mouse CD8a molecule. CD8a is an approximately 32-34 kDa cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha beta) or as a homodimer (CD8 alpha alpha). A majority of thymocytes and a subpopulation of mature alpha beta TCR T cells express CD8 alpha beta while gamma delta TCR T cells, a subpopulation of intestinal intraepithelial lymphocytes (IELs) and dendritic cells express CD8 alpha alpha. CD8 binds to MHC class I and through its association with protein tyrosine kinase p56lck plays a role in T cell development and activation of mature T cells.

Applications Reported

This 53-6.7 antibody has been reported for use in immunohistochemical staining of frozen tissue sections (IHC-F) and immunocytochemistry (ICC).

Applications Tested

This 53-6.7 antibody has been tested by immunohistology of frozen mouse spleen using the IHC/ICC Blocking Buffer - Low Protein (cat. 00-4953). This antibody can be used at 1:100.

For answers to additional questions refer to for IHC/ICC protocols and eFluor Nanocrystal Frequently Asked Questions

Applications: This product has been optimized for use in immunohistochemistry and Immunocytochemistry. We do not recommend its use in flow cytometry. Please refer to cat. 95-0081 as a suitable flow product.

Filter Recommendation: When using this eFluor[®] 650NC antibody conjugate, we recommend a filter that will capture the 650 emission wavelength. A standard Alexa Fluor[®] 647 filter is acceptable, although other filters that narrow the emission would be more suitable when complexing with additional eFluor[®] nanocrystal conjugates. Please refer to Technical Support FAQ for more information.

Buffer Recommendation: We recommend the use of TBS-based solutions when performing IHC/ICC with eFluor[®] NC conjugated antibodies. We offer several products: IHC /ICC Blocking Buffer - Low Protein (cat. 00-4953), and IHC /ICC Blocking Buffer – High Protein (cat. 00-4952) which is optimal when staining FFPE sections or when using eFluor[®] nanocrystal conjugates to nuclear targets.

Mounting Recommendation: For optimal results, we recommend the use of Fluoromount-G[™] (cat. 00-4958) when mounting slides.

References

Mochimaru H, Usui T, Yaguchi T, Nagahama Y, Hasegawa G, Usui Y, Shimmura S, Tsubota K, Amano S, Kawakami Y, Ishida S. Suppression of alkali burn-induced corneal neovascularization by dendritic cell vaccination targeting VEGF receptor 2. Invest Ophthalmol Vis Sci. 2008 May;49(5):2172-7. (53-6.7, in vivo depletion, PubMed)

Yang Z, Day YJ, Toufektsian MC, Xu Y, Ramos SI, Marshall MA, French BA, Linden J. Myocardial infarct-sparing

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com

Anti-Mouse CD8a eFluor® 650NC (for IHC/ICC)

Catalog Number: IH95-0081

Also known as: CD8 alpha, Ly-2, Ly-35, Ly-B, Lyt-2

RUO: For Research Use Only. Not for use in diagnostic procedures.

effect of adenosine A2A receptor activation is due to its action on CD4+ T lymphocytes. *Circulation*. 2006 Nov 7;114(19):2056-64. (53-6.7, in vivo depletion, PubMed)

Taylor JL, Ordway DJ, Troutt J, Gonzalez-Juarrero M, Basaraba RJ, Orme IM. Factors associated with severe granulomatous pneumonia in *Mycobacterium tuberculosis*-infected mice vaccinated therapeutically with hsp65 DNA. *Infect Immun*. 2005 Aug;73(8):5189-93. (53-6.7, IHC frozen)

Grabbe S, Varga G, Beissert S, Steinert M, Pendl G, Seeliger S, Bloch W, Peters T, Schwarz T, Sunderkötter C, Scharffetter-Kochanek K. Beta2 integrins are required for skin homing of primed T cells but not for priming naïve T cells. *J Clin Invest*. 2002 Jan;109(2):183-92. (53-6.7, IHC frozen)

Ledbetter JA, Rouse RV, Micklem HS, Herzenberg LA. T cell subsets defined by expression of Lyt-1,2,3 and Thy-1 antigens. Two-parameter immunofluorescence and cytotoxicity analysis with monoclonal antibodies modifies current views. *J Exp Med*. 1980 Aug 1;152(2):280-95.

Ledbetter, J. A. and L. A. Herzenberg. Xenogeneic monoclonal antibodies to mouse lymphoid differentiation antigens. *Immunol Rev*. 1979;47:63-90.

Related Products

00-4953 IHC /ICC Blocking Buffer - Low Protein

00-4958 Fluoromount-G™

Legal

Under patent number: US 7,939,170 and additional pending patent application(s)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com