

Product Data Sheet

PerCP/Cy5.5 anti-human CD123

Catalog # / Size: 306015 / 25 tests

306016 / 100 tests

Clone: 6H6

Isotype: Mouse IgG1, κ

Immunogen: Human IL-3Rα transfected COS cells Reactivity: Human, Cross-Reactivity: Rhesus

Preparation: The antibody was purified by affinity chromatography, and conjugated with

PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated


PerCP/Cy5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Human peripheral blood lymphocytes stained with 6H6 PerCP/Cy5.5 and **UCHL1 FITC**

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and 564 nm and a maximum emission of 690 nm.

Application Notes: 6H6 antibody does not inhibit IL-3 binding to low- or high-affinity IL-3Rs. Additional reported applications (for the relevant formats) include: Western blotting¹, immunoprecipitation¹, and immunohistochemical staining of acetone-fixed frozen sections2.

> Cy3, Cy5, Cy5.5 and Cy7 are subject to proprietary rights of GE Healthcare Bio-Sciences Corp. and Carnegie Mellon University and made and sold under license from GE Healthcare Bio-Sciences Corp. Sale of this product is licensed

for research use only.

- Application References: 1. Sun Q, et al. 1996. Blood 87:83. (IP, WB)
 - 2. Herling M, et al. 2003. Blood 101:5007. (IHC)
 - 3. Charles N, et al. 2010. Nat. Med. 16:701. (FC) PubMed 4. Martin-Gayo E, et al. 2010. Blood 115:5366. PubMed

Description: CD123 is the 70 kD transmembrane α chain of the IL-3 receptor. Alone, CD123 binds IL-3 with low affinity; when CD123 associates with CDw131 (common β chain), it binds IL-3 with high affinity. CD123 does not transduce intracellular signals upon binding IL-3 and requires the β chain for this function. CD123 is expressed by myeloid precursors, macrophages, dendritic cells, mast cells, basophils, megakaryocytes, and some B cells.

Antigen References: 1. Miyajima A, et al. 1993. Blood 82:1960.

Application Related Products: Product Clone Cell Staining Buffer

FC, ICC, ICFC FC, ICFC FC, ICFC RBC Lysis Buffer (10X)

PerCP/Cy5.5 Mouse IgG1, κ Isotype Ctrl Human TruStain FcX™ (Fc Receptor Blocking Solution) MOPC-21 FC, ICC, ICFC



