

# Product Data Sheet

## LEAF™ Purified anti-human CD129 (IL-9 R)

**Catalog # / Size:** 310408 / 500 µg

**Clone:** AH9R7

**Isotype:** Mouse IgG2b, κ


**Reactivity:** Human

**Preparation:** The LEAF™ (Low Endotoxin, Azide-Free) antibody was purified by affinity chromatography.

**Formulation:** 0.2 µm filtered in phosphate-buffered solution, pH 7.2, containing no preservative. Endotoxin level is <0.1 EU/µg of the protein (<0.01 ng/µg of the protein) as determined by the LAL test.

**Concentration:** 1.0 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C. This LEAF™ solution contains no preservative; handle under aseptic conditions.



Human T lymphoma cell line HUT-78 stained with LEAF™ purified AH9R7, followed by biotinylated anti-mouse IgG and Sav-PE

## Applications:

**Applications:** FC - *Quality tested*  
Block, ELISA - *Reported in the literature*

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.5 µg per 10<sup>6</sup> cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** Additional reported applications (for the relevant formats) include: inhibition the binding of IL-9 to the high affinity α-chain of the human IL-9 receptor<sup>1,2</sup>; and ELISA for detection of soluble IL-9R. For most successful immunofluorescent staining results, it may be important to maximize signal over background by using a relatively bright fluorochrome-antibody conjugate (Cat. No. 310404) or by using a high sensitivity, three-layer staining technique (e.g., including a biotinylated anti-mouse IgG second step (Cat. No. 405303), followed by SAV-PE (Cat. No. 405204)). The LEAF™ Purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 310408).

**Application References:** 1. De Smedt M, *et al.* 2000. *J. Immunol.* 164:1761.  
2. Pilette C, *et al.* 2002. *J. Immunol.* 168:4103.  
3. Demoulin JB, *et al.* 1996. *Mol. Cell. Biol.* 16 (9):4710.

**Description:** CD129 is known as the 57 kD IL-9 receptor. It is a member of the hematopoietin receptor superfamily. Although the α-chain of the receptor binds IL-9 with high affinity, interaction with the γ-chain (CD132) of the IL-2 receptor is required for signaling. The IL-9 receptor is expressed at low levels on eosinophils, mast cells, macrophages, B lymphocytes, T lymphocytes, and erythroid progenitors. IL-9 receptor binding initiates STAT activation required for the proliferative and anti-apoptotic effects of this cytokine. In humans, signals from the IL-9 receptor appear to be critical for intrathymic T cell development. IL-9 binding has been shown to increase IL-5 receptor expression and promote survival in human eosinophils.

**Antigen References:** 1. Demoulin JB, *et al.* 1996. *Mol. Cell. Biol.* 16:4710.

### Related Products:

**Product**  
APC Goat anti-mouse IgG (minimal x-reactivity)  
LEAF™ Purified Mouse IgG2b, κ Isotype Ctrl  
PE Goat anti-mouse IgG (minimal x-reactivity)  
Cell Staining Buffer  
RBC Lysis Buffer (10X)

**Clone**  
Poly4053  
MPC-11  
Poly4053

### Application

FC  
FC, ICFC, WB, IP, ICC, IF, FA  
FC  
FC, ICC, ICFC  
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.