

Product Data Sheet

PE anti-human CD135 (Flt-3/Flk-2)

Catalog # / Size:	313305 / 25 tests 313306 / 100 tests	
Clone:	BV10A4H2	
Isotype:	Mouse IgG1, κ	
Immunogen:	BV-173 pro-B cell line	Relative Cell Number
Reactivity:	Human	
Preparation:	The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.	
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	
Storage:	The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.	10 ⁰ 10 ¹ 10 ² 10 ³ 10 ⁴ Log Fluoresence Intensity
Application	S:	Human pre-B cell line REH stained with BV10A4H2 PE
Applications:	FC - Quality tested	
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. Test size products are transitioning from 20 µl to 5 µl per test . Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 µl staining volume or per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.	
Description:	CD135 is a 130-160 kD type III tyrosine kinase receptor expressed on CD34 ⁺ hematopoietic stem cells, myelomonocytic progenitors, primitive B cell progenitors, and thymocytes. CD135 is also expressed on malignant hematopoietic cells including AML, ALL and CML-BC. CD135, also known as FMS-like tyrosine kinase-3, FLT3, STK-1, and Flk-2, is a growth factor receptor that binds the FLT3 ligand to promote the growth and differentiation of primitive hematopoietic cells. The intracytoplasmic domain of CD135 is modified by phosphorylation and has been shown to interact with Grb2, SOCS1, VAV1, and Shc.	
Antigen References:	1. Rappold I, <i>et al.</i> 1997. <i>Blood</i> 90:111. 2. Rosnet O, <i>et al.</i> 1996. <i>Acta Haematol.</i> 95:218. 3. Rosnet O, <i>et al.</i> 1996. <i>Leukemia</i> 10:238. 4. Bertho JM. <i>et al.</i> 2000. <i>Scand. J. Immunol.</i> 52:53.	

4. Bertho JM, et al. 2000. Scand. J. Immunol. 52:53.

Related Products: Product

PE Mouse IgG1, κ Isotype Ctrl Cell Staining Buffer RBC Lysis Buffer (10X) Human TruStain FcX™ (Fc Receptor Blocking Solution)

Application FC, ICFC FC, ICC, ICFC FC, ICFC FC, ICC, ICFC



Clone

MOPC-21



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.