

Biotin anti-human CD135 (Flt-3/Flk-2)

Catalog # / Size: 313312 / 100 µg

Clone: BV10A4H2

Isotype: Mouse IgG1, κ

Immunogen: BV-173 pro-B cell line


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C. **Do not freeze.**



Human pre-B cell line REH stained with biotinylated BV10A4H2, followed by Sav-PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤2.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Description: CD135 is a 130-160 kD type III tyrosine kinase receptor expressed on CD34⁺ hematopoietic stem cells, myelomonocytic progenitors, primitive B cell progenitors, and thymocytes. CD135 is also expressed on malignant hematopoietic cells including AML, ALL and CML-BC. CD135, also known as FMS-like tyrosine kinase-3, FLT3, STK-1, and Flk-2, is a growth factor receptor that binds the FLT3 ligand to promote the growth and differentiation of primitive hematopoietic cells. The intracytoplasmic domain of CD135 is modified by phosphorylation and has been shown to interact with Grb2, SOCS1, VAV1, and Shc.

Antigen References:

1. Rappold I, *et al.* 1997. *Blood* 90:111.
2. Rosnet O, *et al.* 1996. *Acta Haematol.* 95:218.
3. Rosnet O, *et al.* 1996. *Leukemia* 10:238.
4. Bertho JM, *et al.* 2000. *Scand. J. Immunol.* 52:53.

Related Products:

Product	Clone	Application
Biotin Mouse IgG1, κ Isotype Ctrl	MOPC-21	FC, ICFC
APC Streptavidin		FC, ICFC
APC/Cy7 Streptavidin		FC, ICFC
PE Streptavidin		FC, ICFC
PE/Cy5 Streptavidin		FC, ICFC
PE/Cy7 Streptavidin		FC, ICFC
Cell Staining Buffer		FC, ICC, ICFC
RBC Lysis Buffer (10X)		FC, ICFC
Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.