

## Purified anti-human CD140b (PDGFR $\beta$ )

**Catalog # / Size:** 323602 / 100  $\mu$ g

**Clone:** 18A2

**Isotype:** Mouse IgG1,  $\kappa$

**Immunogen:** NIH-3T3 cells transfected with human PDGFRbeta


**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C.



Human PDGFRB transfected cells stained with purified 18A2, followed by biotinylated anti-mouse IgG and Sav-PE

## Applications:

**Applications:** FC - Quality tested

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is  $\leq 2.0$   $\mu$ g per million cells in 100  $\mu$ l volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** The 18A2 monoclonal antibody recognizes human CD140b also known as the platelet-derived growth factor receptor, beta polypeptide, PDGFR1, and PDGFR $\beta$ . It has been shown to be useful for flow cytometric detection of CD140b.

**Application References:** 1. Vogel W, *et al.* 2002. *Haematologica* 88:126.  
2. Arima, S., *et al.* 2011. *Development*. 138:4763. PubMed.

**Description:** CD140b is a cell surface tyrosine kinase receptor for members of the platelet-derived growth factor family. The identity of the growth factor bound to the receptor determines whether the functional receptor is a homodimer or heterodimer composed of both PDGFR- $\alpha$  and - $\beta$ . CD140b contains two immunoglobulin-like domains and a tyrosine kinase domain with a predicted molecular weight approximately 124 kD. CD140b is widely expressed on a variety of mesenchymal-derived cells and is preferentially expressed on some tumors such as medulloblastoma. Binding of B-chain containing PDGF molecules can stimulate cell proliferation. CD140b has been shown to interact with a number of kinases (including Raf-1, NCK1, FAK, Fyn, others) as well as adaptor molecules and signaling intermediates (Crk, Grb2, Grb4, RasGAP, SHP2, SHC1, others), and has also been shown to associate with integrin  $\beta$ 3 and nexin sorting molecules. CD140b has been implicated in several disease states including atherogenesis and oncogenesis. The PDGFR $\beta$  is heavily phosphorylated on numerous tyrosine residues through both autophosphorylation and ligand-dependent processes.

**Antigen References:** 1. Claesson-Welsh L, *et al.* 1988. *Mol. Cell Biol.* 8:3476.  
2. Gronwald RG, *et al.* 1988. *Proc. Natl. Acad. Sci. USA* 85:3435.  
3. Gilbertson DG, *et al.* 2001. *J. Biol. Chem.* 276:27406.  
4. Seifert RA, *et al.* 1989. *J. Biol. Chem.* 264:8771.  
5. Kanakaraj P, *et al.* 1991. *Biochemistry* 30:1761.

### Related Products: Product

Purified Mouse IgG1,  $\kappa$  Isotype Ctrl  
APC Goat anti-mouse IgG (minimal x-reactivity)  
PE Goat anti-mouse IgG (minimal x-reactivity)  
Cell Staining Buffer

**Clone**  
MOPC-21  
Poly4053  
Poly4053

**Application**  
FC, ICFC, ICC, IF, IHC, IP, WB  
FC  
FC  
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.