

Product Data Sheet

PE anti-human CD134 (OX40)

Catalog # / Size: 350003 / 25 tests

350004 / 100 tests

Clone: Ber-ACT35 (ACT35) **Isotype:** Mouse IgG1, κ

Immunogen: HTLV 1-transformed HUT 102 cells Reactivity: Human, Cross-Reactivity: Chimpanzee

Preparation: The antibody was purified by affinity chromatography and conjugated with PE

under optimal conditions. The solution is free of unconjugated PE and


unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



3-day PHA-activated human peripheral blood lymphocytes were stained with OX-40 (clone Ber-ACT35) PE (filled histogram) or mouse IgG1, κ PE isotype control (open histogram).

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. Test size products are transitioning from 20 µl to 5 µl per test. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 µl staining volume or per 100 µl of whole blood. It is recommended that the reagent be titrated for

optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: Additional reported applications (for the relevant formats) include: Western blotting¹, immunoprecipitation¹, immunohistochemical staining²,³of paraffin embedded⁴ and frozen tissue sections, ELISA⁴, and functional assay⁵. The LEAF™ or Ultra-LEAF™ purified antibody is recommended for functional assays (contact our custom solutions

Application References: 1. Latza U, et al. 1994. Eur. J. Immunol. 24:677. (WB, IP)

Durkop H, et al. 1995. Brit. J. Haematol. 91:927. (IHC)
Durkop H, et al. 1997. Brit. J. Haematol. 98:863. (IHC)
Willett B, et al. 2007. J. Virol. 81:9665. (ELISA)

5. Li M and Zhang Y. et al. 2005. Cell. Mol. Immunol. 2:467. (FA) 6. Gloviczki ML, et al. 2012. Clin J Am Soc Nephrol. PubMed

7. Domingos PL, et al. 2012. An. Bras. Dermatol. 87:851. (IHC)

Description: CD134, also known as OX40 and TNFRSF14, is a 50 kD type I transmembrane glycoprotein. It is a member of the

TNF receptor family. OX40 is expressed on activated T lymphocytes including Th1, Th2, Th17, and Treg cells. The interaction of OX40 with OX40L results in B cell proliferation and antibody secretion, regulation of primary T cell expansion, and T cell survival. OX40 influences the size of the T cell memory pool and regulation of CD4+ T cell

tolerance.

Antigen References: 1. Smith CA, et al. 1994. Cell 76:959. 2. Chen AL, et al. 1999. Immunity 11:689.

3. Croft M. 2010. Annu. Rev. Immunol. 28:57. 4. Ruby CE, et al. 2009. J. Immunol. 183:5079.

5. Klinger M, et al. 2009. J. Immunol. 182:4581.

Related Products: Product Clone Application PE Mouse IgG1, κ Isotype Ctrl (FC) MOPC-21 Cell Staining Buffer FC, ICC, ICFC

RBC Lysis Buffer (10X) FC, ICFC Human TruStain FcX™ (Fc Receptor Blocking Solution) FC, ICC, ICFC



