


Anti-Mouse Granzyme B eFluor® 660 (Alexa Fluor® 647 Replacement)

Catalog Number: 50-8898 Also known as: Gzmb

RUO: For Research Use Only. Not for use in diagnostic procedures.



C57Bl/6 splenocytes were stimulated for 4 days with Anti-Mouse CD3e and Anti-Mouse CD28 Functional Grade Purified (cat. 16-0031 & cat. 16-0281) and Brefeldin A was added for the last 5 hours of culture. Cells were surface stained with Anti-Mouse CD8a FITC (cat. 11-0081) followed by intracellular staining with 0.06 ug of Rat IgG2a K Isotype Control eFluor® 660 (cat. 50-4321) (left) or 0.06 ug of Anti-Mouse Granzyme B eFluor® 660 (right) using IC Fixation & Permeabilization Buffers (cat. 88-8823) and protocol. Total viable cells were used for analysis.

Product Information


Contents: Anti-Mouse Granzyme B eFluor® 660 (Alexa Fluor® 647 Replacement)

Catalog Number: 50-8898

Concentration: 0.2 mg/mL Host/Isotype: Rat IgG2a, kappa



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer **Temperature Limitation:** Store at 2-8°C. Do not



Use By: Refer to vial
Contains sodium azide



Description

This NGZB monoclonal antibody reacts with mouse Granzyme B, which is a member of the granzyme serine protease family. Granzyme B is found in the granules of cytotoxic T cells and NK cells. Granzyme B has also been described as CGL1 (cathepsin G-like-1), a serine protease expressed only in cytotoxic T-lymphocytes after cell activation, and CTLA-1 (cytotoxic T lymphocyte-associated serine esterase 1) based on identification of mRNA in various cytotoxic T cells, but not observed in non-cytotoxic lymphoid cells. Granzyme B is crucial for the rapid induction of target cell death by apoptosis, induced by interaction with cytotoxic T cells. The receptor involved has been identified as mannose 6-phosphate receptor. This receptor functions as a death receptor for Granzyme B during cytotoxic T cell-induced apoptosis. This NGZB monoclonal antibody does not crossreact to human Granzyme B nor is staining blocked with GB11, suggesting it recognizes a different epitope.

Applications Reported

This NGZB antibody has been reported for use in intracellular staining followed by flow cytometric analysis.

Applications Tested

This NGZB antibody has been tested by intracellular staining and flow cytometric analysis of stimulated mouse splenocytes. This can be used at less than or equal to 0.125 μg per test. A test is defined as the amount (μg) of antibody that will stain a cell sample in a final volume of 100 μL . Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

eFluor® 660 is a replacement for Alexa Fluor® 647. eFluor® 660 emits at 659 nm and is excited with the red laser (633 nm). Please make sure that your instrument is capable of detecting this fluorochome.

References

Kapp JA, Honjo K, Kapp LM, Xu XY, Cozier A, Bucy RP. TCR transgenic CD8+ T cells activated in the presence of TGFbeta express FoxP3 and mediate linked suppression of primary immune responses and cardiac allograft rejection. Int Immunol. 2006; 1-14.



An Affymetrix Company

Anti-Mouse Granzyme B eFluor® 660 (Alexa Fluor® 647 Replacement)

Catalog Number: 50-8898 Also known as: Gzmb

RUO: For Research Use Only. Not for use in diagnostic procedures.

Smyth MJ, Trapani JA. Granzymes: exogenous proteinases that induce target cell apoptosis. Immunol Today. 1995 Apr;16(4):202-6.

Related Products

00-4506 Brefeldin A Solution (1000X)
11-0081 Anti-Mouse CD8a FITC (53-6.7)
16-0031 Anti-Mouse CD3e Functional Grade Purified (145-2C11)
16-0281 Anti-Mouse CD28 Functional Grade Purified (37.51)
50-4321 Rat IgG2a K Isotype Control eFluor® 660 (eBR2a)

88-8823 Fixation & Permeabilization Buffers

info@ebioscience.com