

Thermo Scientific DyNAzyme II DNA Polymerase

F-501S/L, 250 U/1000 U

Store at -20°C



1. Introduction

Thermo Scientific DyNAzyme II DNA Polymerase is a thermostable DNA polymerase for routine DNA amplification. It is purified from an *E. coli* strain that carries a plasmid encoding DyNAzyme™ DNA polymerase gene from *Thermus brockianus*. DyNAzyme II DNA polymerase possesses the following activities: 5'→3' DNA polymerase activity and 5'→3' exonuclease activity. DyNAzyme II DNA Polymerase lacks the 3'→5' proofreading activity.

2. Package information

F-501S	250 U (2 U/μl) Material provided: DyNAzyme II DNA Polymerase 250 U (125 μl), 10x Optimized DyNAzyme Buffer (2 x 1.5 ml).
F-501L	1000 U (2 U/μl) Material provided: DyNAzyme II DNA Polymerase 1000 U (500 μl), 10x Optimized DyNAzyme Buffer (4 x 1.5 ml).

Material safety datasheet (MSDS) is available at www.thermoscientific.com/fzmsds.

3. Guidelines for using DyNAzyme II DNA Polymerase

Mix and centrifuge all tubes before opening to ensure homogeneity and improve recovery. Always pipette DyNAzyme II DNA Polymerase carefully and gently. The high glycerol content (50 %) in the storage buffer may otherwise lead to pipetting errors.

Table 1. Pipetting instructions (add items in this order).

Component	Volume	Final conc.
H ₂ O	add to 50 μl	
10x Optimized DyNAzyme buffer	5 μl	1x (1.5 mM MgCl ₂)
10 mM dNTPs	1 μl	200 μM each
Primer A	x μl	0.5 μM*
Primer B	x μl	0.5 μM*
Template DNA	x μl	
DyNAzyme II DNA Polymerase	0.25–1 μl**	0.01–0.04 U/ μM (0.5–2 U/50 μl)

* The recommendation for final primer concentration is 0.5 μM but it can be optimized between 0.2–1.0 μM, if needed.

** Possible enzyme dilutions are recommended to be made in 1x reaction buffer or H₂O immediately before use.

Table 2. Cycling instructions.

Cycle step	2-step protocol		3-step protocol		Cycles
	Temp.	Time	Temp.	Time	
Initial denaturation	94°C	1–2	94°C	1–2 min	1
Denaturation	94°C	15 s–1 min	94°C	15 s–1 min	25–35
Annealing (see 5.3)	–	–	T _m -5°C	10–30 s	
Extension	72°C	40 s/kb	72°C	40 s/kb	
Final extension	72°C 4°C	5–10 min hold	72°C 4°C	5–10 min hold	1

4. Notes about reaction components

4.1 Enzyme

The optimal amount of enzyme depends on the amount of template and the length of the PCR product. Usually 1 unit of DyNAzyme II DNA Polymerase per 50 μl reaction volume gives good results, but the optimal amount can range from 0.5 to 2.0 units per 50 μl reaction depending on the length and difficulty of the amplicon.

4.2 Optimized DyNAzyme Buffer

For standard PCR reactions, optimized DyNAzyme buffer is recommended (1x buffer contains 1.5 mM Mg²⁺). In case Mg²⁺ optimization is required, we recommend DyNAzyme II DNA Polymerase with Mg²⁺-free DyNAzyme buffer and MgCl₂ solution (F-503).

4.3 dNTPs

High quality dNTPs should be used for optimal performance with DyNAzyme II DNA Polymerase. The polymerase also incorporates dUTP, dITP and fluorescently-labeled nucleotides.

4.4 Template

General guidelines for low complexity DNA (e.g. plasmid, lambda or BAC DNA) are: 1 pg–10 ng per 50 μl reaction volume. For high complexity genomic DNA, the amount of DNA template should be 50–500 ng per 50 μl reaction volume. If cDNA synthesis reaction mixture is used directly as a source for the template, the volume used should not exceed 10 % of the final PCR reaction volume.

4.5 PCR additives

PCR additives such as DMSO, formamide, glycerol and betaine are compatible with DyNAzyme II DNA Polymerase. We recommend using PCR additives in the following concentrations: DMSO 2–10 %, formamide 2–10 %, glycerol 5–10 %, or combinations of these. Recommended starting point is 5 % DMSO.

Note: If high DMSO concentration is used, the annealing temperature must be decreased, as DMSO alters the melting point of the primers. It has been reported that 10 % DMSO decreases the annealing temperature by 5.5–6.0°C.¹

5. Notes about cycling conditions

5.1 Denaturation

After an initial 1–2 min denaturation at 94°C, keep the denaturation time as short as possible (usually 30 seconds or less at 94°C).

Note: The denaturation time and temperature also depend on the ramp rate and temperature control mode of the cyclor.

5.2 Primer annealing

The T_m's should be calculated with the nearest neighbor method² as results from primer T_m calculations can vary significantly depending on the method used. Instructions for T_m calculation and a link to a calculator using a modified nearest-neighbor method can be found on www.thermoscientific.com/pcrwebtools. We suggest the primers to be annealed for 30 seconds or less at the highest temperature that will permit annealing of the primers to the template. A guideline for DyNAzyme II DNA Polymerase is to use annealing temperature 5°C below the lower T_m of the primers. Two-step cycling without an annealing step is recommended for high T_m primer pairs.

5.3 Extension

The extension for standard PCR should be performed at 72°C (40 seconds per one kilobase of amplified product).

6. Component specifications

6.1 DyNAzyme II DNA Polymerase (F-501)

The thermostable DyNAzyme II DNA Polymerase is purified from an *E. coli* strain expressing the cloned DyNAzyme DNA Polymerase gene from *Thermus brockianus*, Thermo Scientific's proprietary bacterial strain. DyNAzyme II DNA Polymerase is free of contaminating endo- and exonucleases. It has a half life of 2.5 h at 96°C.

Storage buffer: 20 mM Tris-HCl (pH 7.4 at 25°C), 0.1 mM EDTA, 1 mM DTT, 100 mM KCl, stabilizers, 200 μg/ml BSA and 50 % glycerol.

Unit definition: One unit is defined as the amount of enzyme that will incorporate 10 nmoles of dNTPs into acid-insoluble form at 74°C in 30 minutes under the stated assay conditions.

Unit assay conditions: Incubation buffer: 25 mM TAPS-HCl, pH 9.3 (at 25°C), 50 mM KCl, 2 mM MgCl₂, 1 mM β-mercaptoethanol, 100 μM dCTP, 200 μM each dATP, dGTP, dTTP.

Incubation procedure: 20 μg activated calf thymus DNA and 0.5 μCi [α -³²P] dCTP are incubated with 0.1 units DNA polymerase in 50 μl incubation buffer at 74°C for 10 minutes. The amount of incorporated dNTPs is determined by trichloroacetic acid precipitation.

DNA amplification assay: Performance in PCR is tested by the amplification of 500 bp lambda DNA and 6 kb M13 DNA.

Exonuclease contamination assay: Incubation of 10 U for 4 hours at 72°C in 50 μl assay buffer with 1 μg sonicated [³H] ssDNA (2 x 10⁵ cpm/μg) released <1 % of radioactivity.

Endonuclease contamination assay: No endonuclease activity is observed after incubation of 10 U of DNA polymerase with 1 μg of lambda DNA or lambda HindIII DNA fragments in assay buffer at 72°C for 4 hours.

6.2 10x Optimized DyNAzyme buffer (F-511)

In final 1x reaction concentration the Optimized DyNAzyme buffer contains 10 mM Tris-HCl (pH 8.8 at 25°C), 50 mM KCl, 1.5 mM MgCl₂ and 0.1 % Triton® X-100.

7. References

- Chester N. & Marshak D.R. (1993) *Analytical Biochemistry* 209: 284–290.
- Breslauer K.J. *et al.*, (1986) *PNAS* 83: 3746–3750.

Shipping and storage

DyNAzyme II DNA Polymerase is shipped on gel ice. Upon arrival, store the components at -20°C.

Technical support:

US: techservice.genomics@thermofisher.com

Europe, Asia, Rest of World:

techservice.emea.genomics@thermofisher.com

Web: www.thermoscientific.com/pcr

Tm-calculator: www.thermoscientific.com/pcrwebtools

Product use limitation

This product has been developed and is sold exclusively for research purposes and in vitro use only. This product has not been tested for use in diagnostics or drug development, nor are they suitable for administration to humans or animals.

Designed and manufactured according to certified ISO9001:2008 processes.

© 2011 Thermo Fisher Scientific Inc. All rights reserved. All trademarks are the property of Thermo Fisher Scientific Inc. and its subsidiaries.

v1_07.2011