

Alexa Fluor® 647 anti-human CD160

Catalog # / Size: 341203 / 25 tests
341204 / 100 tests

Clone: BY55


Isotype: Mouse IgM, κ

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 647 under optimal conditions. The solution is free of unconjugated Alexa Fluor® 647.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with BY55 Alexa Fluor® 647 and CD8 (RPA-T8) PerCP/Cy5.5

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is 5 μ l per million cells or 5 μ l per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633nm / 635nm.

** Alexa Fluor® is a registered trademark of Molecular Probes, Inc. Alexa Fluor® dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and issued patents.

Application References:

1. Anumanthan A, *et al.* 1998. *J. Immunol.* 161:2780.
2. Maiza H, *et al.* 1993. *J. Exp. Med.* 178:1121.

Description: CD160 is a 27 kD GPI-anchored glycoprotein also known as BY55, NK1, and NK28. A member the Ig superfamily, CD160 exists as a disulfide-bond multimer, expressed on the surface of a subpopulation of NK cells, γ/δ T cells, subset of CD8+ T cells, and intestinal intraepithelial lymphocytes (IEL). CD160 plays costimulatory roles through binding to classical and nonclassical MHC-I molecules.

Antigen References:

1. Zola H, *et al.* 2007. *Leukocyte and Stromal Cell Molecules: The CD Markers* Wiley-Liss A John Wiley & Sons Inc, Publication.
2. Merino J, *et al.* 2007. *Clin. Exp. Immunol.* 149:87.
3. Barakonyi A, *et al.* 2004. *J. Immunol.* 173:5349.

Related Products:	Product	Clone	Application
	Alexa Fluor® 647 Mouse IgM, κ Isotype Ctrl	MM-30	FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.