

Purified anti-human CD184 (CXCR4)

Catalog # / Size: 306501 / 25 µg
306502 / 100 µg

Clone: 12G5

Isotype: Mouse IgG2a, κ

Workshop Number: VII 70204


Reactivity: Human, **Cross-Reactivity:** Chimpanzee, Baboon, Cynomolgus, Rhesus, Sooty Mangabey, African Green

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Human peripheral blood lymphocytes stained with purified 12G5, followed by biotinylated anti-mouse IgG and Sav-PE

Applications:

Applications: FC - Quality tested
IHC, IF - Reported in the literature
CyTOF® - Validated

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 1.0 µg per 10^6 cells in 100 µl volume or 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections^{2,3}, immunofluorescence microscopy⁶, and blocking of CD4-independent infection by HIV-2 and CD4-dependent infection by some T cell-tropic isolates of HIV-1^{4,5}. Clone 12G5 may not be suitable for Western blotting.¹⁰ The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 306512).

Application References:

- McKnight A, *et al.* 1997. *J. Virol.* 71:1692.
- Endres MJ, *et al.* 1996. *Cell* 87:745. (IHC)
- Volin MV, *et al.* 1998. *Biochem. Biophys. Res. Commun.* 242:46. (IHC)
- Berndt C, *et al.* 1998. *P. Natl. Acad. Sci. USA* 95:12556. (Block)
- Ullrich CK, *et al.* 2000. *Blood* 96:1438. (Block)
- Murga M, *et al.* 2005. *Blood* 105:1992. (IF)
- Thompson BD. 2007. *J. Biol. Chem.* 282:9547. (FC) PubMed
- Isnardi I, *et al.* 2010. *Blood* 115:5026. PubMed
- Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
- Fischer T, *et al.* 2008. *PLoS One* 3:e4069.

Description: CD184, also known as fusin or CXCR4, is a 45 kD seven transmembrane G-protein-linked CXC chemokine receptor. CD184 is widely expressed on blood and tissue cells, including B and T cells, monocytes, macrophages, dendritic cells, granulocytes, megakaryocytes/platelets, lymphoid, myeloid precursor cells, endothelial cells, epithelial cells, astrocytes, and neurons, among other tissue cells. CD184 is the receptor for CXC chemokine SDF-1, mediates blood cell migration, and is involved in B lymphopoiesis and myelopoiesis, cardiogenesis, blood vessel formation, and cerebellar development. CXCR4 is also a coreceptor of X4 HIV-1 and an alternative receptor for some isolates of HIV-2. The 12G5 antibody has been reported to block CD4-independent infection by HIV-2 and CD4-dependent infection by some T cell-tropic isolates of HIV-1.

Antigen References:

- Berger E, *et al.* 1999. *Annu. Rev. Immunol.* 17:657.
- Loetscher P, *et al.* 2000. *Adv. Immunol.* 74:127.
- Murphy P, *et al.* 2000. *Pharmacol. Rev.* 52:145.

Related Products:	Product	Clone	Application
	APC Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC
	Purified Mouse IgG2a, κ Isotype Ctrl	MOPC-173	FC, ICC, IF, IHC, IP, WB
	PE Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	Purified anti-human CD184 (CXCR4)	4G10	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.