

Product Data Sheet

PE anti-human CD184 (CXCR4)

Catalog # / Size: 306505 / 25 tests
306506 / 100 tests

Clone: 12G5

Isotype: Mouse IgG2a, κ


Workshop Number: VII 70204

Reactivity: Human, **Cross-Reactivity:** Chimpanzee, Baboon, Cynomolgus, Rhesus, Sooty Mangabey, African Green

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with 12G5 PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections^{2,3}, immunofluorescence microscopy⁶, and blocking of CD4-independent infection by HIV-2 and CD4-dependent infection by some T cell-tropic isolates of HIV-1^{4,5}. Clone 12G5 may not be suitable for Western blotting.¹⁰ The LEAF™ purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 306512).

Application References:

- McKnight A, *et al.* 1997. *J. Virol.* 71:1692.
- Endres MJ, *et al.* 1996. *Cell* 87:745. (IHC)
- Volin MV, *et al.* 1998. *Biochem. Biophys. Res. Commun.* 242:46. (IHC)
- Berndt C, *et al.* 1998. *P. Natl. Acad. Sci. USA* 95:12556. (Block)
- Ullrich CK, *et al.* 2000. *Blood* 96:1438. (Block)
- Murga M, *et al.* 2005. *Blood* 105:1992. (IF)
- Thompson BD. 2007. *J. Biol. Chem.* 282:9547. (FC) PubMed
- Isnardi I, *et al.* 2010. *Blood* 115:5026. PubMed
- Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
- Fischer T, *et al.* 2008. *PLoS One* 3:e4069.

Description: CD184, also known as fusin or CXCR4, is a 45 kD seven transmembrane G-protein-linked CXC chemokine receptor. CD184 is widely expressed on blood and tissue cells, including B and T cells, monocytes, macrophages, dendritic cells, granulocytes, megakaryocytes/platelets, lymphoid, myeloid precursor cells, endothelial cells, epithelial cells, astrocytes, and neurons, among other tissue cells. CD184 is the receptor for CXC chemokine SDF-1, mediates blood cell migration, and is involved in B lymphopoiesis and myelopoiesis, cardiogenesis, blood vessel formation, and cerebellar development. CXCR4 is also a coreceptor of X4 HIV-1 and an alternative receptor for some isolates of HIV-2. The 12G5 antibody has been reported to block CD4-independent infection by HIV-2 and CD4-dependent infection by some T cell-tropic isolates of HIV-1.

Antigen References:

- Berger E, *et al.* 1999. *Annu. Rev. Immunol.* 17:657.
- Loetscher P, *et al.* 2000. *Adv. Immunol.* 74:127.
- Murphy P, *et al.* 2000. *Pharmacol. Rev.* 52:145.

Related Products:

Product
PE Mouse IgG2a, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)
PE anti-human CD184 (CXCR4)
Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone
MOPC-173

4G10

Application
FC, ICFC
FC, ICC, ICFC
FC, ICFC
FC
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.