

APC/Cy7 anti-human CD206 (MMR)

Catalog # / Size: 321119 / 25 tests
321120 / 100 tests

Clone: 15-2

Isotype: Mouse IgG1, κ


Immunogen: Purified human mannose receptor

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC/Cy7 under optimal conditions. The solution is free of unconjugated APC/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



GM-CSF- stimulated (day-3) human monocytes stained with 15-2 APC/Cy7

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: The 15-2 antibody blocks the interaction of MMR with its ligand, and inhibits mannose receptor-mediated degradation of t-PA by macrophages. Additional reported applications of this antibody (for the relevant formats) include: Western blotting¹, blocking of ligand binding^{1,2}, immunofluorescence³, and immunohistochemical staining of acetone-fixed frozen tissue sections¹. The LEAF™ purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 321112).

Cy3, Cy5, Cy5.5 and Cy7 are subject to proprietary rights of GE Healthcare Bio-Sciences Corp. and Carnegie Mellon University and made and sold under license from GE Healthcare Bio-Sciences Corp. Sale of this product is licensed for research use only.

Application References:

- Noorman F, *et al.* 1997. *J. Leukocyte Biol.* 61:63. (WB, IHC, Block)
- Barrett-Bergshoeff M, *et al.* 1997. *Thromb Haemost.* 77:718. (Block)
- Kato M, *et al.* 2007. *J. Immunol.* 179:6052. (IF)
- Isik G, *et al.* 2013. *PLoS One.* 8:60126. PubMed.

Description: Macrophage mannose receptor (MMR) is a 162-175 kD type I membrane protein also known as CD206, MRC1, or mannose receptor (MR). It is a pattern recognition receptor (PRR) that belongs to C-type lectin superfamily. MMR is expressed on macrophages, dendritic cells, and hepatic or lymphatic endothelial cells, but not on monocytes. MMR recognizes a range of microbial carbohydrates bearing mannose, fucose, or N-acetyl glucosamine. MMR mediates endocytosis and phagocytosis, induces activation of macrophages and antigen presentation, plays an important role in host defense, and provides a link between innate and adaptive immunity.

Antigen References:

- Mason D, *et al.* Eds. 2002. *Leukocyte Typing VII.* Oxford University Press. p303
- Wileman TE, *et al.* 1986. *P. Natl. Acad. Sci. USA* 83:2501.
- Apostolopoulos V and McKenzie IF. 2001. *Curr. Mol. Med.* 1:469.
- Le Cabec V, *et al.* 2005. *J. Leukocyte Biol.* 77:934.
- Barrett-Bergshoeff M, *et al.* 1997. *Thromb. Haemostatis* 77:718.

Related Products:	Product	Clone	Application
	APC/Cy7 Mouse IgG1, κ Isotype Ctrl	MOPC-21	FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.