

## Biotin anti-human CD202b (Tie2/Tek)

**Catalog # / Size:** 334203 / 25 µg  
334204 / 100 µg

**Clone:** 33.1 (Ab33)

**Isotype:** Mouse IgG1 κ

**Immunogen:** Recombinant extracellular domain of human Tie2


**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C. **Do not freeze.**



Human endothelial cell line, HUVEC, stained with biotinylated 33.1, followed by Sav-PE

## Applications:

**Applications:** FC -Quality tested

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 2.0 µg per 10<sup>6</sup> cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** Additional reported applications include: immunoprecipitation, Western blot, immunohistochemical staining of frozen tissue sections and ELISA

**Application References:** 1. Peters KG, *et al.* 1998. *Br. J. Cancer.* 77:51.  
2. Wong AL, *et al.* 1997. *Circ. Res.* 81:567.  
3. Lin P, *et al.* 1998. *P. Natl. Acad. Sci. USA* 95:8829.

**Description:** CD202b is a 145 kD type I transmembrane protein, also known as Tie2 or TEK. It is a member of the receptor tyrosine kinase (RTK) family of proteins and is expressed by endothelial cells and their progenitors, quiescent hematopoietic stem cells (HSCs), and a subpopulation of monocytes. Angiopoietin-1 (Ang-1) is an activator of CD202b to promote, maintain, and stabilize mature vessels and to maintain HSCs in quiescent state. Ang-2 is another ligand of CD202b, which is involved in postnatal angiogenesis and in antagonizing the effects of Ang-1. Tie2 binds to Ang-4 as well.

**Antigen References:** 1. De Palma M, *et al.* 2005. *Cancer Cell.* 8:211  
2. Shaw JP, *et al.* 2004. *Blood Cells Mol. Dis.* 32:168  
3. Hsu HC, *et al.* 2000. *Blood.* 96:3757  
4. Arai F, *et al.* 2004. *Cell.* 118:149

### Related Products:Product

Cell Staining Buffer  
Human TNF-α ELISA MAX™ Standard  
Biotin Mouse IgG1, κ Isotype Ctrl  
Human TruStain FcX™ (Fc Receptor Blocking Solution)

### Clone

MOPC-21

### Application

FC, ICC, ICFC  
ELISA  
FC, ICFC  
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.