

Mouse IL-5 Recombinant Protein

Catalog Number: 14-8051 Also Known As:Interleukin-5, IL5 RUO: For Research Use Only

Product Information

Contents: Mouse IL-5 Recombinant Protein

REF Catalog Number: 14-8051

Handling Conditions: For best recovery, quick-spin vial prior to

opening. Use in a sterile environment

Source: Insect cells infected with baculovirus: amino acids Met

21- Gly 133 of mouse IL-5 (accession # NM_010558).

Molecular Mass: The protein has a predicted molecular mass of 13,126. The DTT reduced protein migrates as two major bands of 16 and 21 kDa and the non-reduced protein migrates as a group of bands corresponding to dimer at 37-39 kDa. Heterogeneity results from different glyscosylation states. Purity: Greater than 98%, as determined by SDS-PAGE

Endotoxin Level: Less than 0.01 ng/ug cytokine as determined

by the LAL assay.

Bioactivity: Measured by human TF-1 cell proliferation assay. The ED_{50} is 0.1 ng/ml, corresponding to a specific activity of 1.0 x 10^7 Units/mg.

Formulation: Sterile liquid; 10 mM sodium phosphate, pH 7.2, 150 mM NaCl, 0.5% BSA. 0.22 µm filtered.

150 mivi Naci, 0.5% BSA. 0.22 μm filtered.

Temperature Limitation: Store at less than or equal to -70°C.

Batch Code: Refer to Vial
Use By: Refer to Vial

Description

Recombinant mouse IL-5 is a disulfide-linked homodimer, containing two 113 amino acid peptides; these resolve as 32-34 kDa bands in SDS-PAGE. IL-5 is produced by T cells and has been known as eosinophil-differentiating factor (EDF), B cell growth factor II (BCGFII), and T cell-replacing factor (TRF). IL-5 induces eosinophil differentiation and promotes eosinophil survival and activation. In mice, IL-5 has been shown to stimulate B cell proliferation and antibody production.

Applications Reported

Recombinant mouse IL-5 is biologically active and is useful for bioassays and as a cytokine standard in mouse IL-5 ELISA

Applications Tested

This recombinant mouse IL-5 has been tested by sandwich ELISA using the anti-mouse IL-5 antibody pair (TRFK5/TRFK4), and in bioassays using the IL-5 responsive human cell line TF-1. The ED₅₀ measured in a TF-1 proliferation assay is typically 100 pg/ml, corresponding to a specific activity of approximately 1×10^7 Units/mg.

Related Products

00-4202 ELISA Diluent Solution (5X)

13-7051 Anti-Mouse IL-5 Biotin (TRFK4)

14-7052 Anti-Human/Mouse IL-5 Purified (TRFK5)

18-4100 Avidin HRP

44-2404 Nunc MaxiSorp® flat-bottom 96 well plate

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com