

Human IL-27 Recombinant Protein

Catalog Number: 14-8279

Also Known As: Interleukin-27, EBI3, p29, IL-30, IL27

RUO: For Research Use Only

Product Information

Contents: Human IL-27 Recombinant Protein

REF **Catalog Number:** 14-8279

Handling Conditions: For best recovery, quick-spin vial prior to opening. Use in a sterile environment

Source: Human EBI3 (met 1-lys229; accession #NM_005755) linked through a (gly-gly-gly-gly-ser)₃ to mature P28/IL-30 (phe 29-pro 243; accession # NM_145659) tagged at the C-terminal with thr-gly-his₁₀ was expressed in human 293 cells.


Molecular Mass: After removal of the secretion signal the mature EBI3-linker-mature P28-His₁₀ protein has a predicted molecular mass of approximately 50,000. On non-reducing and reducing SDS-PAGE the linked protein migrates as a 55 kDa protein due to glycosylation.

Purity: Greater than 98%, as determined by SDS-PAGE


Endotoxin Level: Less than 0.01 ng/ug cytokine, as determined by the LAL assay.

Bioactivity: Measured in a bioassay of IL-27-mediated inhibition of IL-2 production by mouse splenocytes activated with immobilized anti-CD3 and soluble anti-CD28 antibodies (Villarino, A.V., et al. 2006. J. Immunol. 176: 237). The ED₅₀ for this is typically below 100 ng/ml, corresponding to a specific activity of greater than 1.0 x 10⁴ U/mg.

Formulation: Sterile liquid; 20 mM NaH₂PO₄, pH 6.0, 0.6M NaCl, with 1.0% BSA. 0.22 µm filtered.

 **Temperature Limitation:** Store at less than or equal to -70°C.

LOT **Batch Code:** Refer to Vial

 **Use By:** Refer to Vial

Description

IL-27, a member of the IL-12 family, is a heterodimeric protein consisting of the p40-related protein Epstein-Barr virus-induced gene 3 (EBI3) non-covalently linked to an IL-12p35-related protein, p28 (also known as IL-30). IL-27 is produced by activated APCs and mature dendritic cells. IL-27 exerts its activities on NK cells and naïve CD4⁺ T cells; mRNA expression analysis of IL-27 receptor components (WSX-1/TCCR and gp130) suggests that IL-27 may also target other cells, including mast cells and monocytes. Binding of IL-27 to WSX-1/gp130 activates JAK1, STAT1, and STAT3 and STAT1/3 phosphorylation. WSX-1/TCCR-deficient mice develop impaired Th1 responses and are more susceptible to infection with *L. monocytogenes* suggesting that Th1 responses require IL-27. Although activation of WSX-1 is required for the initiation of Th1 responses, it is not necessary for maintaining Th1 responses. IL-27 alone is not able to induce the differentiation of CD4⁺ T cells into IFN-γ-producing cells, suggesting a role for IL-27 as an initial activator of Th1 responses. An important effect of IL-27 in initiating Th1 responses is the induction of the Th1-specific transcription factor T-bet as well as the suppression of the Th2-specific transcription factor GATA-3. T-bet plays a critical role in Th1 differentiation by its ability to maintain IL-12Rβ₂ expression following CD4⁺ T cell activation.

Recent studies indicate that IL-27 has a potent antitumor activity. In vitro, IL-27 has been found to act directly on naïve CD8 cells, generating CTL with enhanced granzyme B expression. In vivo, IL-27 has been reported to augment CTL activity, inhibit tumor growth, and induce complete regression of primary and metastatic neuroblastoma tumors.

Applications Reported

The recombinant mouse IL-27 (EBI3/p28) has been reported useful for bioassay.

Applications Tested

This recombinant human IL-27 (EBI3/p28) has been tested in bioassay for inhibition of IL-2 production by mouse splenocytes activated with immobilized anti-CD3 and soluble anti-CD28 antibodies (Villarino, A.V., et al. 2006. J. Immunol. 176: 237). The ED₅₀ for this is typically below 100 ng/ml, corresponding to a specific activity of greater than 1.0 x 10⁴ U/mg.

References

Stumhofer, J.S., et al. 2006. IL-27 negatively regulates the development of IL-17-producing T helper cells during chronic inflammation of the central nervous system. *Nature Immunol.* 7: 937-45.

Villarino, A., et al. 2006. IL-27 Limits IL-2 Production during Th1 Differentiation. J. Immunol. 176: 237 - 247.
Salcedo, R., et al. 2004. IL-27 mediates complete regression of orthotopic primary and metastatic murine neuroblastoma tumors: role for CD8+ T cells. J. Immunol. 173: 7170-7182.
Morishima, N., et al. 2005. Augmentation of effector CD8+ T cell generation with enhanced granzyme B expression by IL-27. J. Immunol. 175: 1686-1693.
Owaki, T., et al. 2005. A role for IL-27 in early regulation of Th1 Differentiation. J. Immunol. 175: 2191-2200.

Related Products

14-8129 Human IL-12 p70 Recombinant Protein
14-8239 Human IL-23 Recombinant Protein
14-8348 Human TGF beta 1 Recombinant Protein
88-7239 Human IL-23 (p19/p40) Platinum ELISA
88-7344 Human/Mouse TGF beta 1 ELISA Ready-SET-Go![®] (To Be Discontinued. Refer to 2nd Generation RSG Version: cat. 88-8350)

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com