


Human TGF beta 3 Recombinant Protein

Catalog Number: 14-8369

Also known as: Transforming Growth Factor beta 3, TGF-beta3, TGFbeta3

RUO: For Research Use Only. Not for use in diagnostic procedures.



Human TGF beta3 Recombinant Protein neutralization of recombinant Mouse IL-4-induced proliferation in CTLL-2 cells.

Product Information

Contents: Human TGF beta 3 Recombinant Protein

REF **Catalog Number:** 14-8369

Concentration: 0.1 mg/mL

Handling Conditions: For best recovery, quick-spin vial prior to opening. Use in a sterile environment

Source: E. coli derived, accession number NM_003239

Molecular Mass: 25.7 kDa

Purity: > 98%, as determined by SDS-PAGE

Endotoxin: Less than 0.01 ng/ug cytokine as determined by the LAL assay

Bioactivity: The ED50 of this protein, as measured by the neutralization of recombinant mouse IL-4-induced proliferation of CTLL-2 cells, is 0.03-0.125 ng/mL. This corresponds to a specific activity of $3.3 \times 10^7 - 8 \times 10^6$ Units/mg.

Formulation: Sterile liquid: 0.1 M glycine, pH 3.0 with 1% BSA, 0.22 um filtered

Temperature Limitation: Store at less than or equal to -70°C.

Batch Code: Refer to vial

Use By: Refer to vial



Description

Transforming growth factor $\beta 3$ (TGF- $\beta 3$) is the third member of the transforming growth factor family of cytokines, which also includes TGF- $\beta 1$ and - $\beta 2$. These cytokines are secreted in precursor form consisting of a bioactive C-terminal domain attached to an N-terminal domain known as latency associated protein (LAP). Cleavage of LAP results in the mature protein, which functions as a disulfide-linked homodimer. As with all members of the family, TGF $\beta 3$ is highly conserved across species, with mouse and human TGF beta 3 demonstrating 100% sequence homology and cross-species activity.

The members of this family can be expressed by most cell types and exert pleiotropic effects, which include the suppression of B- and T-cell effector activity, mediation of tissue healing, and suppression of tumor proliferation. The promotion of CD4+CD25+ T-cell expansion is a newly discovered function of the TGF- β cytokines, and indicates an important role in the protection against autoimmunity.

Applications Reported

Recombinant human TGF- $\beta 3$ is biologically active.

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com

Human TGF beta 3 Recombinant Protein

Catalog Number: 14-8369

Also known as: Transforming Growth Factor beta 3, TGF-beta3, TGFbeta3

RUO: For Research Use Only. Not for use in diagnostic procedures.

Applications Tested

The ED₅₀ of this protein, as measured by the neutralization of recombinant mouse IL-4-induced proliferation of CTLL-2 cells, is 0.03-0.125 ng/ml. This corresponds to a specific activity of 3.3×10^7 - 8×10^6 Units/mg.

References

Cox DA and Maurer T. Transforming Growth Factor- β . Clin Immunol. 1997 Apr;83(1)25-30.

Peng Y, Laouar Y, Li MO, Green EA, Flavell RA. TGF-beta regulates in vivo expansion of Foxp3-expressing CD4+CD25+ regulatory T cells responsible for protection against diabetes. Proc Natl Acad Sci U S A. 2004 Mar 30;101(13):4572-7.

Zheng SG, Gray JD, Ohtsuka K, Yamagiwa S, Horwitz DA. Generation ex vivo of TGF-beta-producing regulatory T cells from CD4+CD25- precursors. J Immunol. 2002 Oct 15;169(8)4183-9.

Related Products

14-8348 Human TGF beta 1 Recombinant Protein

14-8368 Human TGF beta 2 Recombinant Protein

88-7344 Human/Mouse TGF beta 1 ELISA Ready-SET-Go!® (To Be Discontinued. Refer to 2nd Generation RSG Version: cat. 88-8350)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com