


## Human IL-3 Recombinant Protein Carrier-Free

Catalog Number: 34-8039

Also Known As: Interleukin-3, IL3

RUO: For Research Use Only. Not for use in diagnostic procedures.



Proliferation of the TF-1 cell line in response to Human IL-3 Recombinant Protein Carrier-Free

### Product Information

Contents: Human IL-3 Recombinant Protein Carrier-Free

**REF** Catalog Number: 34-8039

Handling Conditions: For best recovery, quick-spin vial prior to opening. Use in sterile environment.

Source: *E. coli* expressed mature human IL-3 (accession #NM\_000588).


Molecular Mass: 15 kDa

Purity: Greater than 95%, as determined by SDS-PAGE.


Endotoxin Level: Less than 0.01 ng/ug cytokine as determined by the LAL assay.

Bioactivity: The ED<sub>50</sub> measured in a TF-1 proliferation assay is typically less than 0.15 ng/ml, corresponding to a specific activity of greater than 7x10<sup>6</sup> Units/mg.

Formulation: Sterile liquid: 20mM phosphate buffer, 0.15M NaCl, pH 7.0, 0.22 µm filtered.

 Temperature Limitation: Store at less than or equal to -70°C.

**LOT** Batch Code: Refer to Vial

 Use By: Refer to Vial

### Description

Interleukin 3 (IL-3), also called Mast Cell Growth Factor (MCGF), Multi-Colony Stimulating Factor (multi-CSF), Eosinophil-CSF (E-CSF), is produced by activated T cells, mast cells and eosinophils. IL-3 is a hematopoietic growth factor that stimulates colony formation of erythroid, megakaryocyte, neutrophil, eosinophil, basophil, mast cell and monocytic lineages. Most of these functions are enhanced or dependent on co-stimulation with other cytokines. Human and mouse IL-3 share only 29% sequence identity at the amino acid level. The bioactivity of IL-3 is highly species specific. Human IL-3 has no activity on mouse cells.

### Applications Reported

Recombinant human IL-3 is biologically active.

### Applications Tested

This recombinant human IL-3 has been tested by bioassays using the TF-1 cell line. The ED<sub>50</sub> measured in a TF-1 proliferation assay is typically less than 0.15 ng/ml, corresponding to a specific activity of greater than 7x10<sup>6</sup> Units/mg.

### References

Robin C, Ottersbach K, Durand C, Peeters M, Vanes L, Tybulewicz V, Dzierzak E. An unexpected role for IL-3 in the embryonic development of hematopoietic stem cells. *Dev Cell*. 2006 Aug;11(2):171-80.

Celestin J, Rotschke O, Falk K, Ramesh N, Jabara H, Strominger J, Geha RS. IL-3 induces B7.2 (CD86) expression and costimulatory activity in human eosinophils. *J Immunol*. 2001 Dec 1;167(11):6097-104.

Donahue RE, Seehra J, Metzger M, Lefebvre D, Rock B, Carbone S, Nathan DG, Garnick M, Sehgal PK, Laston D. Human IL-3 and GM-CSF act synergistically in stimulating hematopoiesis in primates. *Science*. 1988 Sep 30;241(4874):1820-3.

Yang YC, Ciarletta AB, Temple PA, Chung MP, Kovacic S, Witek-Giannotti JS, Leary AC, Kriz R, Donahue RE, Wong GG. Human IL-3 (multi-CSF): identification by expression cloning of a novel hematopoietic growth factor related to murine IL-3. *Cell*. 1986 Oct 10;47(1):3-10.

Related Products  
14-8031 Mouse IL-3 Recombinant Protein

---

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.eBioscience.com](http://www.eBioscience.com) • [info@eBioscience.com](mailto:info@eBioscience.com)