

Product Data Sheet

PE anti-human Perforin

Catalog # / Size: 308105 / 25 tests
308106 / 100 tests

Clone: dG9

Isotype: Mouse IgG2b, κ


Immunogen: Purified granules from the human lymphoma cell line

Reactivity: Human, **Cross-Reactivity:** Bovine (Cow, Cattle); does not cross-react with mouse¹.

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes intracellularly stained with dG9 PE

Applications:

Applications: ICFC - *Quality tested*

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation, intracellular flow cytometric analysis and immunofluorescence microscopy^{5,7}, and immunohistochemical staining of acetone-fixed frozen tissue sections and formalin-fixed paraffin-embedded tissue sections^{1,4}.

Application References:

1. Hameed A, et al. 1992. *Am. J. Pathol.* 140:1025. (IHC)
2. Schaeferli P, et al. 2004. *J. Exp. Med.* 199:1265.
3. Watanabe N, et al. 1997. *Blood* 90:3662.
4. Mauad T, et al. 2004. *Pediatr. Pulmonol.* 38:233. (IHC)
5. Barrat FJ, et al. 1999. *P. Natl. Acad. Sci. USA* 96:8645. (IF)
6. Chen H, et al. 2005. *J. Immunol.* 175:591.
7. Bryceson YT, et al. 2007. *Blood* doi:10.1182/blood-2007-02-074468. (IF)
8. Wood SM, et al. 2009. *Blood* 114:4117. PubMed

Description: Perforin is a 70 kD cytolytic protein that is expressed in the cytoplasmic granules of cytotoxic T lymphocytes (CTLs) and natural killer (NK) cells. Perforin is one of the major effector molecules used by cytotoxic T cells and NK cells to mediate targeted cell lysis.

Antigen References:

1. Lieberman J. 2003. *Nat. Rev. Immunol.* 3:361.
2. Trapani J, et al. 2002. *Nat. Rev. Immunol.* 2:735.

Related Products:

Product
PE Mouse IgG2b, κ Isotype Ctrl

Cell Staining Buffer

RBC Lysis Buffer (10X)

PE anti-human Granzyme A

Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone
MPC-11

CB9

Application

FC, ICFC

FC, ICC, ICFC

FC, ICFC

ICFC

FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.