

Purified anti-Phosphotyrosine

Catalog # / Size: 309301 / 25 µg
309302 / 100 µg

Clone: PY20

Isotype: Mouse IgG2b, κ

Immunogen: KLH-conjugated phosphotyrosine


Reactivity: All Species

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 50% glycerol. Final antibody concentration is 0.5 mg/ml.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored at -20°C.



Hela cell extract was resolved by electrophoresis, transferred to nitrocellulose, and probed with monoclonal anti-phosphotyrosine antibody (clone PY-20). Lane 1, serum-starved Hela cells; Lane 2, serum-starved Hela cells following serum addition for 4 hrs. Lane 2 shows an upregulation of tyrosine phosphorylated proteins after serum addition. Proteins were visualized using a goat anti-mouse secondary conjugated to HRP and a chemiluminescence detection system.

Applications:

Applications: WB - *Quality tested*
ICFC, IF, IP- *Reported in the literature*

Recommended Usage: Western blotting, suggested working dilution(s): Use 1-2 µg per ml antibody dilution buffer per mini-gel. Do not use dilution or blocking buffers containing milk as they may interfere with antibody binding to proteins of interest. Dilution and blocking buffers containing 4% bovine serum albumin are recommended for use with this antibody. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation^{1,2}, Western blotting^{1,2}, immunofluorescence microscopy³.

Application References:

1. Vuori K, *et al.* 1995. *J. Biol. Chem.* 270:22259. (IP, WB)
2. Glenney J, *et al.* 1988. *J. Immunol. Meth.* 109:277. (IP, WB)
3. Prahalad P, *et al.* 2004. *Am J Physiol Cell Physiol* 286:C693. (IF)
4. Zentillin L, *et al.* 2009. *FASEB J.* 24:1467. PubMed
5. Prochazka R, *et al.* 2012. *Reproduction.* 144:535. PubMed.

Description: Phosphorylation is a common modification of proteins that can result in alterations in protein function, protein-protein association, cellular localization, and protein-half life. Phosphorylation can occur on threonine, serine, and tyrosine residues. The PY20 monoclonal antibody recognizes phosphorylated tyrosine residues in all species tested (human, mouse, rat, dog, chicken, and frog). The PY20 antibody has been shown to be useful for flow cytometry, immunoprecipitation, Western blotting, and immunofluorescence staining.

Related Products:	Product	Clone	Application
	Purified anti-mouse CD3	17A2	FC, IHC, IP
	Purified anti-human CD3	HIT3a	FC, IHC, IP
	Purified anti-human CD3	UCHT1	FC, IHC, IP, WB, CyTOF®
	Purified anti-Fyn	Poly6041	WB
	Purified anti-Lyn	Poly6046	WB
	HRP Donkey anti-rabbit IgG (minimal x-reactivity)	Poly4064	ELISA, IHC, WB



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.