

Biotin anti-human CD354 (TREM-1)

Catalog # / Size: 314904 / 100 µg

Clone: TREM-26

Isotype: Mouse IgG1, κ

Immunogen: Recombinant human TREM-1


Reactivity: Human, **Cross-Reactivity:** Dog (Canine)

Preparation: The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C. **Do not freeze.**



Applications:

Applications: FC - *Quality tested*
ELISA Detection - *Validated in house*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.5 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: **ELISA Detection:** The biotinylated TREM-26 antibody is useful as a detection antibody for a TREM-1 specific sandwich ELISA, when used in conjunction with the TREM-37 antibody (Cat. No. 316102) as the capture antibody. The TREM-26 antibody is useful for flow cytometry and activation of monocytes and granulocytes. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 314907).

Application References: 1. Li J, *et al.* 2011. *Dev. Comp. Immunol.* epub.

Description: TREM-1 is a 30 kD glycoprotein also known as triggering receptor expressed on myeloid cells 1, and triggering receptor expressed on monocytes 1. It is a Type I membrane protein that contains an immunoglobulin-like V-type domain. Alternatively spliced protein variant may be secreted. TREM-1 is highly expressed on peripheral blood myeloid cells (particularly mature monocytes and granulocytes); TREM-1 expression can be further upregulated by bacteria, fungi and lipopolysaccharide. TREM-1 has been shown to interact with the adaptor protein DAP12 to stimulate neutrophil and monocyte-mediated inflammatory responses through the triggering and release of pro-inflammatory cytokines and chemokines. TREM-1 is thought to amplify inflammatory responses to fungal and bacterial infections and potentiate septic shock. This antibody has been shown to be useful for flow cytometry and activation of monocytes and granulocytes.

Antigen References: 1. Bouchon A, *et al.* 2001. *Nature* 410:1103.
2. Bouchon A, *et al.* 2000. *J. Immunol.* 164:4991.
3. Gingras MC, *et al.* 2002. *Mol. Immunol.* 38:817.

Related Products:	Product	Clone	Application
	Biotin Mouse IgG1, κ Isotype Ctrl	MOPC-21	FC, ICFC
	PE Streptavidin		FC, ICFC
	Cell Staining Buffer		FC, ICC, ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.