

## **Product Data Sheet**

## APC anti-human Ig light chain $\lambda$

Catalog # / Size: 316609 / 25 tests

316610 / 100 tests

Clone: MHL-38

**Isotype:** Mouse IgG2a, κ **Immunogen:** Human Ig cocktail

Reactivity: Human

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with

APC under optimal conditions. The solution is free of unconjugated APC and


unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Human peripheral blood lymphocytes stained with CD19 PE and MHL-38

## **Applications:**

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. Test size products are transitioning from 20 μl to 5 μl per test. Please check your vial or your CoA to find the

size products are transitioning from 20 μl to 5 μl per test. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μl staining volume or per 100 μl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at

www.biolegend.com/testsize regarding the test size change.

Description: The MHL-38 antibody reacts with both soluble and membrane human immunoglobulin light chain lambda (λ). It does

not react with human immunoglobulin light chain kappa (κ) or heavy chain (μ). The MHL-38 antibody can be used as

primary or secondary reagent for immunofluorescent staining or ELISA analysis.

Related Products: Product

APC Mouse IgG2a, κ Isotype Ctrl

Cell Staining Buffer

Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone MOPC-173 Application FC, ICFC FC, ICC, ICFC FC, ICC, ICFC



