

Purified anti-human CD23

Catalog # / Size: 317002 / 100 µg

Clone: D3.6

Isotype: Mouse IgG2a, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Human peripheral blood lymphocytes stained with purified D3.6, followed by anti-mouse IgG FITC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.5 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Description: CD23 is a 45 kD protein, also known as Leu-20, FcεRII, IgE Fc receptor, BLAST-2, B6, and low affinity IgE receptor. It is a member of the Ig family, expressed on most mature B cells, B cells in follicular mantle (but not in proliferating germinal center cells, follicular dendritic cells, monocytes, eosinophils, Langerhans cells, and a subset of T cells (10-15% of tonsillar T cells). CD23 responds to high levels of IgE by downregulating IgE secretion. In human monocytes, CD23 triggering results in release of pro-inflammatory cytokines including TNF-α, IL-1, IL-6, and GM-CSF. CD23 can be proteolytically cleaved to generate soluble CD23 fragments of various molecular weights. In chronic lymphocytic leukemia, levels of soluble CD23 in the serum can be used as a prognostic marker to identify patients at high risk for disease progression. Alternate splicing of exon 2 can also generate two cell-surface isoforms of CD23 differing by 6 amino acids in their cytoplasmic region. The D3.6 antibody against human CD23 has been shown to be useful for flow cytometry.

Antigen References:

1. Ludin C, *et al.* 1987. *EMBO J.* 6:109.
2. Delespesse G, *et al.* 1992. *Immunol. Rev.* 125:77.
3. Flores-Romo L, *et al.* 1993. *Science* 261:1038.
4. Armant M, *et al.* 1994. *J. Exp. Med.* 180:1005.

Related Products: Product

APC Goat anti-mouse IgG (minimal x-reactivity)
Biotin Goat anti-mouse IgG (minimal x-reactivity)
Purified Mouse IgG2a, κ Isotype Ctrl
PE Goat anti-mouse IgG (minimal x-reactivity)
Cell Staining Buffer

Clone

Poly4053
Poly4053
MOPC-173
Poly4053

Application

FC
FC, ELISA, IHC, IF, WB
FC, ICC, IF, IHC, IP, WB
FC
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.