

Product Data Sheet

Biotin anti-Cytochrome c

Catalog # / Size: 612303 / 100 µg

Clone: 6H2.B4

Isotype: Mouse IgG1, κ

Immunogen: Rat cyt c-OVA

Reactivity: Mouse, Rat, Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C. **Do not freeze.**

Applications:

Applications: ICFC - *Quality tested*
IP, IF, ICC

Recommended Usage: For immunofluorescent staining, the suggested use of this reagent is ≤ 0.5 µg per 10^6 cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: intracellular flow cytometry⁵, immunofluorescence microscopy^{3,5}, immunoprecipitation⁴, and immunocytochemistry⁵.

Application References:

1. Goshorn SC, *et al.* 1991. *J. Biol. Chem.* 266:2134.
2. Jemmerson R, *et al.* 1991. *Eur. J. Immunol.* 21:143.
3. Chandra D, *et al.* 2002. *J. Biol. Chem.* 277:50842. (IF)
4. Semenkova L, *et al.* 2003. *Eur. J. Biochem.* 270:4388. (IP)
5. Shih S-F, *et al.* 2001. *J. Biol. Chem.* 276:21870. (ICFC ICC IF)

Description: Cytochrome c is a 15 kD protein found in the mitochondrial intermembrane space with a heme-binding domain. Cytochrome c is a component of the electron transport chain; the heme group transfers electrons from cytochrome b-c1 complex to cytochrome oxidase complex. Cytochrome c initiates apoptosis by release to cytoplasm and binding Apaf-1 which activates procaspase 9. Cytochrome c interacts with the cytochrome b-c1 complex, cytochrome oxidase complex, heme, Apaf-1, and Caspase 9 proteins. The 6H2.B4 monoclonal antibody recognizes human, mouse, and rat cytochrome-c and has been shown to be useful for intracellular flow cytometric staining, Western blotting, immunoprecipitation, and immunofluorescence staining.

Antigen References:

1. Liu X, *et al.* 1996. *Cell.* 86:147.
2. Li P, *et al.* 1997. *Cell.* 91:479.
3. Zhang Z, *et al.* 2003. *Gene* 312:61.
4. Ferguson H, *et al.* 2003. *J. Biol. Chem.* 278:45793.

Related Products:

Product
Biotin Mouse IgG1, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)

Clone
MOPC-21

Application
FC, ICFC
FC, ICC, ICFC
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.