

# Product Data Sheet

## Recombinant Mouse FGF-basic (carrier-free)

**Catalog # / Size:** 579602 / 10 µg  
579604 / 25 µg  
579606 / 100 µg  
579608 / 500 µg

**Source:** Mouse FGF-basic, amino acids Ala11-Ser154 (Accession# NM\_008006), was expressed in *E. coli*.

**Molecular Mass:** The 144 amino acid recombinant protein has a predicted molecular mass of 16,223 Da. The DTT-reduced and non-reduced protein migrate at approximately 16kDa by SDS-PAGE. The N-terminal amino acid is Alanine.

**Purity:** Purity is >98%, as determined by Coomassie stained SDS-PAGE.

**Endotoxin Level:** Endotoxin level is <0.1 EU/µg (<0.01ng/µg) protein as determined by the LAL method.

**Activity:** ED50 = 0.3 to 2 ng/ml, corresponding to a specific activity of 3.3 to 0.5 x 10<sup>6</sup> units/mg, as determined by the dose dependent stimulation of 3T3 cell proliferation.

**Preparation:** 10-100µg sizes are bottled at 200µg/mL. 500µg sizes and larger are bottled at the concentration indicated on the vial.

**Formulation:** 0.22 µm filtered protein solution is in 10mM NaH<sub>2</sub>PO<sub>4</sub>, 150mM NaCl, pH 7.2, 1mM DTT, 0.5mM EDTA.

**Storage:** Unopened vial can be stored at -20°C for six months or at -70°C for one year. For maximum results, quick spin vial prior to opening. Stock solutions should be prepared at no less than 10µg/mL in buffer containing carrier protein such as 1% BSA or HSA or 10% FBS. For long term storage, aliquot into polypropylene vials and store in a manual defrost freezer. **Avoid repeated freeze/thaw cycles.**

## Applications:

**Applications:** Bioassay

**Description:** FGFb is a member of the fibroblast growth factor (FGF) family which includes 23 members. FGFb is expressed in almost all tissues and play important roles in a variety of normal and pathological processes, including development, wound healing and neoplastic transformation. FGFb is mitogenic for many cell types, both epithelial and mesenchymal. FGFb shows potent angiogenic activity and has been implicated in tumor angiogenesis. FGF-2 significantly promoted the proliferation of adipose-derived mesenchymal cells (AMC) and enhanced chondrogenesis in three-dimensional micromass culture. FGFb binds to a family of four distinct, high affinity tyrosine kinase receptors, designated FGFR-1 to -4. In addition, FGFb binds to the ECM, and heparan sulfate (HS) is an essential and dynamic regulator of fibroblast growth factor (FGF) signaling. Two fundamentally different crystallographic models have been proposed to explain, at the molecular level, how HS/heparin enables FGF and FGF receptor (FGFR) to assemble into a functional dimer on the cell surface, although there is controversy regarding the exact manner by which this occurs.

- Antigen References:**
1. Rusnati M and Presta M *Current Pharm Des* 13:2025-2044 2007.
  2. Chaffer CL, *et al. Differentiation* 75(9):831-42 2007.
  3. Cronauer NV, *et al. Eur Urol* 43:309-319 2003.
  4. Shimizu A, *et al. J Biol Chem* 276:11031-11040 2001.
  5. Mahammadi M, *et al. Curr Opin Struct Biol* 15:506-516 2005.
  6. Chiou M Xu Y Longaker MT *Biochem Biophys Res Commun* 343:644-652 2006.



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.