

# Product Data Sheet

## APC/Cy7 anti-human IL-17A

**Catalog # / Size:** 512319 / 25 tests  
512320 / 100 tests

**Clone:** BL168

**Isotype:** Mouse IgG1,  $\kappa$


**Immunogen:** Recombinant full length human IL-17A

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography and conjugated with APC/Cy7 under optimal conditions. The solution is free of unconjugated APC/Cy7 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Storage:** The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



PMA + ionomycin-stimulated (6 hours) human peripheral blood lymphocytes surface stained with CD3 FITC and then intracellularly stained with BL168 APC/Cy7 (top) or mouse IgG1,  $\kappa$  APC/Cy7 isotype control (bottom)

## Applications:

**Applications:** ICFC - *Quality tested*

**Recommended Usage:** Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20  $\mu$ l to 5  $\mu$ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100  $\mu$ l staining volume or per 100  $\mu$ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at [www.biolegend.com/testsize](http://www.biolegend.com/testsize) regarding the test size change.

Cy3, Cy5, Cy5.5 and Cy7 are subject to proprietary rights of GE Healthcare Bio-Sciences Corp. and Carnegie Mellon University and made and sold under license from GE Healthcare Bio-Sciences Corp. Sale of this product is licensed for research use only.

**Description:** IL-17A is the founding member of the IL-17 family, a group of six structurally related pro-inflammatory cytokines. IL-17A, secreted by activated CD4<sup>+</sup> Th17 cell subpopulation, elicits multiple biological activities on a variety of cells including: the induction of IL-6, IL-8, G-CSF, and PGE2 production in epithelial, endothelial or fibroblasts; the enhancement of surface expression of ICAM-1 in fibroblasts; activation of NF- $\kappa$ B and costimulation of T cell proliferation. Recent studies demonstrated that, in mice, activated IL-17-secreting CD4<sup>+</sup> helper T cells (Th17 cells) mediate an autoimmune arthritis that clinically and immunologically resembles rheumatoid arthritis (RA). Human IL-17A shows 63%, 63%, and 72% amino acid sequence identity to rat IL-17A, mouse IL-17A, and a protein encoded by the ORF13 gene of herpesvirus Saimiri (HVS), respectively.

**Antigen References:**


1. Hirota K, *et al.* 2007. *J. Exp. Med.* 204:41.
2. Furuzawa-Carballeda J, *et al.* 2007. *Autoimmun. Rev.* 6:169.
3. Witowski J, *et al.* 2007. *Kidney Int.* 71:514.
4. Gaffen SL, *et al.* 2006. *Vitam. Horm.* 74:255.
5. Hymowitz S, *et al.* 2001. *EMBO J.* 20:5332.

### Related Products:

**Product**  
APC/Cy7 Mouse IgG1,  $\kappa$  Isotype Ctrl  
Cell Staining Buffer  
RBC Lysis Buffer (10X)

**Clone**  
MOPC-21

**Application**  
FC, ICFC  
FC, ICC, ICFC  
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.