

Product Data Sheet

FITC anti-human IL-8

Catalog # / Size: 514604 / 100 tests

Clone: BH0814

Isotype: Mouse IgG2b, λ


Immunogen: Recombinant human IL-8, amino acids Ser28-Ser99 (Accession # NM_000584)

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



LPS-stimulated (6 hours, in the presence of monensin) human peripheral blood monocytes intracellularly stained with BH0814 FITC

Applications:

Applications: ICFC - *Quality tested*

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: ELISA Capture: The purified BH0814 antibody is useful as the capture antibody in a sandwich ELISA assay, when used in conjunction with the biotinylated BH0840 (Cat. No. 514703/ 514704) antibody as the detecting antibody and recombinant human IL-8 (Cat. # 570909) as the standard.

Description: IL-8, also known as neutrophil chemotactic factor, neutrophil activating protein, and monocyte-derived neutrophil chemotactic factor, is a member of the alpha (C-X-C) subfamily of chemokines called CXCL8. In response to proinflammatory stimuli, IL-8 is produced by monocytes, macrophages, T cells, neutrophils, and fibroblasts. IL-8 promotes neutrophil chemotaxis and degranulation. The 72 amino acid IL-8 is the predominant form secreted by monocytes and lymphocytes.

Antigen References:

1. Fitzgerald K, *et al.* Eds. 2001. The Cytokine FactsBook. Academic Press San Diego.
2. Baggiolini M, *et al.* 1994. *Adv. Immunol.* 55:97.
3. Schröder J, *et al.* 1992. *Immunology Ser.* 57:387.
4. Zwahlen R, *et al.* 1993. *Intl. Rev. Expt. Pathol.* 34:27.

Related Products:

Product
 Cell Staining Buffer
 RBC Lysis Buffer (10X)
 Fixation Buffer
 Permeabilization Wash Buffer (10X)
 Brefeldin A Solution (1,000X)
 Monensin Solution (1,000X)

Clone

Application

FC, ICC, ICFC
 FC, ICFC
 ICC, ICFC
 ICC, ICFC, IHC
 ICFC
 ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.