

LEAF™ Purified anti-human IL-9

Catalog # / Size: 507704 / 500 µg

Clone: MH9A3

Isotype: Mouse IgG1, κ

Immunogen: Baculovirus-expressed, recombinant human IL-9

Reactivity: Human

Preparation: The LEAF™ (Low Endotoxin, Azide-Free) antibody was purified by affinity chromatography.

Formulation: 0.2 µm filtered in phosphate-buffered solution, pH 7.2, containing no preservative. Endotoxin level is <0.1 EU/µg of the protein (<0.01 ng/µg of the protein) as determined by the LAL test.

Concentration: 1.0 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C. This LEAF™ solution contains no preservative; handle under aseptic conditions.

Applications:

Applications: ELISA - *Quality tested*
Neut - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by ELISA assay. For use as an ELISA detection antibody, a concentration range of 0.5-2 µg/ml is recommended. To obtain a linear standard curve, serial dilutions of IL-9 recombinant protein ranging from 1000 to 8 pg/ml are recommended for each ELISA plate. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: **ELISA Detection⁴:** The biotinylated MH9A3 antibody is useful as a detection antibody for a human IL-9 sandwich ELISA assay, when used in conjunction with Purified MH9A4 antibody (Cat. No. 507602) as the capture antibody. **Neutralization^{2,3}:** The LEAF™Purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for neutralization of human IL-9 bioactivity (Cat. No. 507704).

Application References:

1. Jenmalm M, *et al.* 2001. *Clin. Exptl. Aller.* 31:1528.
2. Personal communication with developer.
3. Gounni AS, *et al.* 2004. *J. Immunol.* 173:2771.
4. Faulkner H, *et al.* 2002. *J. Infec. Diseases.* 185:665.

Description: IL-9 is a potent, T cell-derived, T cell growth factor which can also enhance mast cell activity and IL-3- or IL-4-dependent proliferation of bone marrow-derived mast cells. IL-9 synergizes with erythropoietin to promote erythroid colony formation. IL-9 has also been reported to protect human T cells from apoptosis induced by IL-2 withdrawal. IL-9 is upregulated in human eosinophils by TNF-α and IL1-β. IL-9 has been reported to downregulate the oxidative burst in activated human alveolar macrophages and induce TGF-β production. The MH9A3 antibody reacts with human IL-9. The MH9A3 antibody can inhibit IL-9 bioactivity *in vitro*.

Antigen References:

1. Fitzgerald K, *et al.* Eds. 2001. *The Cytokine FactsBook.* Academic Press San Diego.
2. Quesniaux V. 1992. *Research Immunology* 143:385.
3. Renaud J, *et al.* 1993. *Adv. Immunol.* 54:79.
4. Yang Y. 1992. *Leuk. Lymphoma* 8:441.

Related Products:Product

LEAF™ Purified Mouse IgG1, κ Isotype Ctrl
Recombinant Human IL-9

Clone
MOPC-21

Application
FC, ICFC, WB, IP, ICC, IF, FA
BA, ELISA



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.