

Product Data Sheet

Purified anti-human IL-8

Catalog # / Size: 511402 / 500 µg

Clone: E8N1

Isotype: Mouse IgG1, κ

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.

Applications:

Applications: ELISA

Recommended Usage: Each lot of this antibody is quality control tested by ELISA assay. For ELISA detection applications, a concentration range of 0.25-1.0 µg/ml is recommended. To obtain a linear standard curve, serial dilutions of IL-8 recombinant protein ranging from 2000 to 15 pg/ml are recommended for each ELISA plate. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: **ELISA Detection:** The biotinylated E8N1 antibody is useful as the detection antibody in a sandwich ELISA assay, when used in conjunction with the purified H8A5 (Cat. No. 511502) antibody as the capture antibody.
Note: For testing human IL-8 in serum or plasma, BioLegend's ELISA Max™ Sets are specially developed and recommended.

Description: IL-8, also known as neutrophil chemotactic factor, neutrophil activating protein, and monocyte-derived neutrophil chemotactic factor, is a member of the alpha (C-X-C) subfamily of chemokines called CXCL8. In response to proinflammatory stimuli, IL-8 is produced by monocytes, macrophages, T cells, neutrophils, and fibroblasts. IL-8 promotes neutrophil chemotaxis and degranulation. The 72 amino acid IL-8 is the predominant form secreted by monocytes and lymphocytes. The E8N1 antibody recognizes the human IL-8 protein and has been shown to be useful for intracellular immunofluorescence flow cytometric analysis and as ELISA detection antibody.

Antigen References: 1. Fitzgerald K, *et al.* Eds. 2001. The Cytokine FactsBook. Academic Press San Diego.
2. Baggiolini M, *et al.* 1994. *Adv. Immunol.* 55:97.
3. Schröder J, *et al.* 1992. *Immunology Ser.* 57:387.
4. Zwahlen R, *et al.* 1993. *Intl. Rev. Expt. Pathol.* 34:27.

Related Products:

Product
Purified Mouse IgG1, κ Isotype Ctrl
Cell Staining Buffer
Fixation Buffer
Permeabilization Wash Buffer (10X)

Clone
MOPC-21

Application
FC, ICFC, ICC, IF, IHC, IP, WB
FC, ICC, ICFC
ICC, ICFC
ICC, ICFC, IHC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.