

Product Data Sheet

FITC anti-mouse IFN- γ

Catalog # / Size: 505805 / 25 μ g
505806 / 100 μ g

Clone: XMG1.2

Isotype: Rat IgG1, κ

Immunogen: *E. coli*-expressed, recombinant mouse IFN- γ


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The IFN- γ antibody solution should be stored undiluted at 4°C, and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 mouse splenocytes were stimulated with PMA + Ionomycin for 6 hours (in the presence of monensin), stained with CD3 APC, fixed, permeabilized, and then stained with IFN- γ (clone XMG1.2) FITC (top) or rat IgG1, κ FITC isotype control (bottom).

Applications:

Applications: ICFC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 1.0 μ g per 10^6 cells in 100 μ l volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: **ELISA^{1-4,11,14} or ELISPOT⁵ Detection:** The biotinylated XMG1.2 antibody is useful as a detection antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with purified R4-6A2 antibody (Cat. No. 505702/505706) as the capture antibody and recombinant mouse IFN- γ (Cat. No. 575309) as the standard.

ELISA or ELISPOT Capture: The purified XMG1.2 antibody is useful as a capture antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with biotinylated R4-6A2 antibody (Cat. No. 505704) as the detection antibody and recombinant mouse IFN- γ (Cat. No. 575309) as the standard. The LEAFTM purified antibody is suggested for ELISPOT capture (Cat. No. 505812).


Flow Cytometry^{7,8,12,13,16}: The fluorochrome-labeled XMG1.2 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IFN- γ -producing cells within mixed cell populations.

Neutralization^{1-3,9,10}: The XMG1.2 antibody can neutralize the bioactivity of natural or recombinant IFN- γ . The LEAFTM purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for neutralization of mouse IFN- γ bioactivity *in vivo* and *in vitro* (Cat. No. 505812). For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAFTM purified antibody (Cat. No. 505834) with a lower endotoxin limit than standard LEAFTM purified antibodies (Endotoxin <0.01 EU/ μ g).

Additional reported applications (for the relevant formats) include:

Western blotting, immunohistochemical staining of paraformaldehyde-fixed, saponin-treated frozen tissue sections⁶, and immunocytochemistry.

Note: For testing mouse IFN- γ in serum, plasma or supernatant, BioLegend's ELISA MaxTM Sets (Cat. No. 430801 to 430806) are specially developed and recommended.



- Application References:**
1. Abrams J, *et al.* 1992. *Immunol. Rev.* 127:5. (ELISA, Neut)
 2. Sander B, *et al.* 1993. *J. Immunol. Meth.* 166:201. (ELISA, Neut)
 3. Abrams J, *et al.* 1995. *Curr. Prot. Immunol.* John Wiley and Sons, New York. Unit 6.20. (ELISA, Neut)
 4. Yang X, *et al.* 1993. *J. Immunoassay* 14:129. (ELISA)
 5. Klinman D, *et al.* 1994. *Curr. Prot. Immunol.* John Wiley and Sons, New York. Unit 6.19. (ELISPOT)
 6. Sander B, *et al.* 1991. *Immunol. Rev.* 119:65. (IHC)
 7. Ferrick D, *et al.* 1995. *Nature* 373:255. (FC)
 8. Ko SY, *et al.* 2005. *J. Immunol.* 175:3309. (FC) PubMed
 9. Peterson KE, *et al.* 2000. *J. Virol.* 74:5363. (Neut)
 10. DeKrey GK, *et al.* 1998. *Infect. Immun.* 66:827. (Neut)
 11. Dzhalalov I, *et al.* 2007. *J. Immunol.* 178:2113. (ELISA)



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

12. Lawson BR, *et al.* 2007. *J. Immunol.* 178:5366. (FC)
13. Lee JW, *et al.* 2006. *Nature Immunol.* 8:181. (FC) PubMed
14. Xu G, *et al.* 2007. *J. Immunol.* 179:5358. (ELISA) PubMed
15. Montfort M, *et al.* 2004. *J. Immunol.* 173:4084. PubMed
16. Haring JS, *et al.* 2008. *J. Immunol.* 180:2855. (FC) PubMed
17. Jordan JM, *et al.* 2008. *Infect Immun.* 76:3717. PubMed
18. Tonkin DR, *et al.* 2008. *J. Immunol.* 181:4516. PubMed
19. Charles N, *et al.* 2010. *Nat. Med.* 16:701. (FC) PubMed
20. Cui Y, *et al.* 2009. *Invest. Ophthalm. Vis. Sci.* 50:5811. (FC) PubMed

Description: Interferon- γ is a potent multifunctional cytokine which is secreted primarily by activated NK cells and T cells. Originally characterized based on anti-viral activities, IFN- γ also exerts anti-proliferative, immunoregulatory, and proinflammatory activities. IFN- γ can upregulate MHC class I and II antigen expression by antigen-presenting cells.

Antigen References: 1. Fitzgerald K, *et al.* Eds. 2001. The Cytokine FactsBook. Academic Press, San Diego.
 2. De Maeyer E, *et al.* 1992. *Curr. Opin. Immunol.* 4:321.
 3. Farrar M, *et al.* 1993. *Annu. Rev. Immunol.* 11:571.
 4. Gray P, *et al.* 1987. *Lymphokines* 13:151.

Related Products:	Product	Clone	Application
	Cell Staining Buffer		FC, ICC, ICFC
	Fixation Buffer		ICC, ICFC
	Permeabilization Wash Buffer (10X)		ICC, ICFC, IHC
	Brefeldin A Solution (1,000X)		ICFC
	Monensin Solution (1,000X)		ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	FITC Rat IgG1, κ Isotype Ctrl	RTK2071	FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.