

Product Data Sheet

PE anti-mouse/human IL-5

Catalog # / Size: 504303 / 25 µg
504304 / 100 µg

Clone: TRFK5

Isotype: Rat IgG1, κ

Immunogen: Partially-purified T cell clone supernatant

Reactivity: Mouse, Human, **Cross-Reactivity*:** Guinea Pig

Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**

Applications:

Applications: ICFC - *Quality tested*

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: **ELISA^{1,3,4} or ELISPOT^{2,6} Capture:** The purified TRFK5 antibody is useful as a capture antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with biotinylated TRFK4 antibody (Cat. No. 504402) as the detection antibody for mouse IL-5, or used in conjunction with biotinylated JES1-5A10 antibody (Cat. No. 501002/501005) as the detection antibody for human IL-5. The LEAF™ purified antibody is suggested for ELISPOT capture.

Flow Cytometry⁵: The fluorochrome-labeled TRFK5 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IL-5-producing cells within mixed cell populations. For intracellular cytokine staining protocol, please visit www.biolegend.com and click on the support section.

Neutralization^{1,9}: The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for neutralization of mouse and human IL-5 bioactivity (Cat. No. 504308).

Additional reported applications (for the relevant formats) include: Western blotting, immunohistochemical staining of formalin-fixed paraffin-embedded tissue sections⁸ and paraformaldehyde-fixed, saponin-treated frozen tissue sections⁷, and immunocytochemistry.

Note: For testing human IL-5 in serum or plasma, BioLegend's ELISA Max™ Sets (Cat. No. 430401 to 430406) are specially developed and recommended. For testing mouse IL-5 in serum, plasma or supernatant, BioLegend's ELISA Max™ Sets (Cat. No. 431201 to 431206) are specially developed and recommended.

Application References:

1. Abrams, J. 1995. *Curr. Prot. Immunol.* John Wiley and Sons, New York. Unit 6.20.
2. Klinman, D., et al. 1994. *Curr. Prot. Immunol.* John Wiley and Sons, New York. Unit 6.19.
3. Swain, S., et al. 1990. *J. Immunol.* 145:3796.
4. Abrams, J., et al. 1992. *Immunol. Rev.* 127:5.
5. Assenmacher, M., et al. 1994. *Eur. J. Immunol.* 24:1097.
6. Karulin, A., et al. 2000. *J. Immunol.* 164:1862.
7. Andersson, U., et al. 1999. *Detection and quantification of gene expression.* New York:Springer-Verlag.
8. Fan, W. Y., et al. 2001. *Exp. Biol. Med.* 226:1045.
9. Sitkauskienė, B., et al. 2005. *Respir Res.* 6:33.

Description: IL-5 is a homodimeric, disulphide-linked protein produced by T-cells. Monomeric human IL-5 is a 126 amino acid protein with a reported molecular weight of 26 kD for the homodimeric protein. Mouse and human IL-5 are approximately 70% identical. IL-5 has been shown to promote the growth of immature hematopoietic BFU-E progenitors, stimulate the activation and differentiation of eosinophils, and promote the generation of cytotoxic lymphocytes. Mouse IL-5 induces the differentiation and proliferation of pre-activated B-cells and stimulates the production and secretion of IgM and IgA by B-cells stimulated with bacterial endotoxin. The TRFK5 antibody can neutralize the bioactivity of natural or recombinant IL-5.

Antigen References:

1. Fitzgerald, K., et al. Eds. 2001. *The Cytokine FactsBook.* Academic Press, San Diego.
2. Takatsu, K., et al. 1988. *Immunol. Rev.* 102:107.
3. Takatsu, K. 1992. *Curr. Opin. Immunol.* 4:299.
4. Takatsu, K. 1991. *Microbiol. Immunol.* 35:593.

Related Products: Product
Cell Staining Buffer

Clone

Application
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

Fixation Buffer
Permeabilization Wash Buffer (10X)
Brefeldin A Solution (1,000X)
Monensin Solution (1,000X)
RBC Lysis Buffer (10X)
PE Rat IgG1, κ Isotype Ctrl

RTK2071

ICC, ICFC
ICC, ICFC, IHC
ICFC
ICFC
FC, ICFC
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible
for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 9727 Pacific Heights Blvd, San Diego, CA 92121 www.biollegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587