

Product Data Sheet

Purified anti-mouse IL-10

Catalog # / Size: 505001 / 50 µg

505002 / 500 µg

Clone: JES5-16E3 **Isotype:** Rat IgG2b, κ

Immunogen: E. coli-expressed, recombinant mouse IL-10


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Mouse IL-10 Concentration (pg/mL)

Applications:

Applications: ELISA Capture - Quality tested

IHC - Reported in the literature

CyTOF® - Validated

Recommended Usage: Each lot of this antibody is quality control tested by ELISA assay. For ELISA capture applications, a concentration

range of 0.1-1.0 µg/ml is recommended. It is recommended that the reagent be titrated for optimal performance for each application. To obtain a linear standard curve, serial dilutions of IL-10 recombinant protein ranging from 2000 to

15 pg/ml are recommended for each ELISA plate. For ELISA capture, Carbonate buffer, pH 9.5 is highly recommended for coating buffer.

Application Notes: ELISA or ELISPOT Detection^{1,9,11}: The biotinylated JES5-16E3 antibody is useful as a detection antibody for a

sandwich ELISA or ELISPOT assay, when used in conjunction with purified JES5-2A5 antibody (Cat. No.

504902/504904) as the capture antibody.

Neutralization¹⁴: The LEAF™ Purified JES5-16E3 antibody can neutralize the bioactivity of natural or recombinant

IL-10.

Application References:

Simkin G, et al. 2000. J. Immunol. 164:2457.
Kitagaki K, et al. 2002. Clin. Diagn. Lab Immunol. 9:1260.

3. Khanna A, et al. 2000. J. Immunol. 164:1346.

Sander B, et al. 1993. J. Immunol. Methods 166:201.
Litton M, et al. 1994. J. Immunol. Methods 175:47.

6. Andersson U, et al. 1999. Detection and qunatification of gene expression. New York: Springer-Verlag.

7. Finkelman F, *et al.* 2003. *Curr. Prot. Immunol.* John Wiley & Sons New York. Unit 6.28. 8. Wang W, *et al.* 2004. *FASEB J.* 18:1043. 9. Brummel R and Lenert P. 2005. *J. Immunol.* 174:2429. 10. Lawson BR, et al. 2007. J. Immunol. 178:5366. 11. Xu G, *et al.* 2007. *J. Immunol.* 179:5358. PubMed 12. Brummel R, *et al.* 2005. *J. Immunol.*174:2429. PubMed 13. Kang YJ, et al. 2007. Stem Cells 25:1814. PubMed

14. Seo N, et al. 2001. Immunology. 103:449. (Neut)

Description: IL-10 was originally described as Cytokine Synthesis Inhibitory Factor (CSIF) by virtue of its ability to inhibit cytokine

production by Th1 clones. IL-10 shares over 80% sequence homology with the Epstein-Barr virus protein BCRFI. IL-10 inhibits IFN- γ , TNF- β , and IL-2 production by Th1 clones; inhibits macrophage-mediated IL-1, IL-6, and TNF- α synthesis; suppresses the delayed type hypersensitivity response; stimulates Th2 cell response (which results in

elevated antibody production); and promotes mast cell proliferation in combination with IL-4.

Antigen References: 1. Fitzgerald K, et al. Eds. 2001. The Cytokine FactsBook. Academic Press San Diego.

2. de Waal-Malefy R, et al. 1992. Curr. Opin. Immunol. 4:314. 3. Howard M, et al. 1992. Immunol. Today 13:198.

 Quesniaux V. 1992. Res. Immunol. 143:385. 5. Norton SK, et al. 2008. J. Immunol. 180:2848.

Related Products: Product Application Clone

Recombinant Mouse IL-10 rm IL-10 BA, ELISA HRP Avidin ELÍSA, ELISPOT, IHC, WB Avidin TMB Substrate Reagent Set **ELISA** ELISA Assay Diluent (5X) **ELISA**

Biotin anti-mouse IL-10 JES5-2A5 ELISA Detection, ELISPOT



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible on for patent infringement or other violations that may occur with the use of our products.





