

Purified anti-human MCP-1

Catalog # / Size: 502601 / 50 µg
502602 / 500 µg

Clone: 5D3-F7

Isotype: Mouse IgG1, κ

Immunogen: Recombinant human MCP-1

Reactivity: Human, **Cross-Reactivity*:** Cynomolgus, Rhesus

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.

Applications:

Applications: ELISA Capture - *Quality tested*
ELISA & ELISPOT Detection, ICFC, IHC, IP, WB - *Reported in the literature*

Recommended Usage: Recommended Usage: Each lot of this antibody is quality control tested by ELISA assay. For ELISA capture applications, a concentration range of 2-8 µg/ml is recommended. To obtain a linear standard curve, serial dilutions of MCP-1 recombinant protein ranging from 2000 to 31.3 pg/ml are recommended for each ELISA plate. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: **ELISA or ELISPOT Detection¹:** The biotinylated 5D3-F7 antibody is useful as the detection antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the purified 2H5 antibody (Cat. No. 505902/505906) as the capture antibody.
ELISA or ELISPOT Capture: The purified 5D3-F7 antibody is useful as the capture antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the biotinylated 2H5 antibody (Cat. No. 505908) as the detection antibody. The LEAF™ purified antibody (Cat. No. 502607) is suggested for ELISPOT capture.
Additional reported applications (for the relevant formats) include: intracellular flow cytometry², immunoprecipitation^{1,3}, Western blotting¹, and immunohistochemical staining¹.

Application References:

1. Peri, G., *et al.* 1994. *J. Immunol. Meth.* 174:249.
2. Rezaie-Majd, A., *et al.* 2002. *Arterioscler Thromb Vasc Biol.* 22:1194.
3. Hirsch, A., *et al.* 1999. *J. Virol.* 73:404.

Description: Monocyte chemoattractant protein-1 (MCP-1) also known as monocyte chemoattractant and activating factor (MCAF) was identified based on its ability to chemoattract monocytes. Subsequently, MCP-1 has also been found to regulate adhesion molecule expression and cytokine production in monocytes. MCP-1 is identical to the product of the JE gene, a PDGF inducible gene. MCP-1 is a member of the beta (C-C) chemokine subfamily, known as CCL2. The 5D3-F7 antibody reacts with human monocyte chemoattractant protein-1 (MCP-1).

Antigen References:

1. Fitzgerald, K., *et al.* Eds. 2001. *The Cytokine FactsBook.* Academic Press, San Diego.
2. Bischoff, S., *et al.* 1992. *J. Exp. Med.* 175:1271.
3. Charo, I., *et al.* 1994. *P. Natl. Acad. Sci. USA* 91:2752.
4. Jiang, Y., *et al.* 1992. *J. Immunol.* 148:2423.

Related Products:	Product	Clone	Application
	Purified anti-mouse/rat/human MCP-1	2H5	ELISA Capture, IHC, WB



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.