

## APC anti-mouse IL-2

**Catalog # / Size:** 503809 / 25 µg  
503810 / 100 µg

**Clone:** JES6-5H4

**Isotype:** Rat IgG2b, κ

**Immunogen:** *E. coli*-expressed, recombinant mouse IL-2


**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.2 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



PMA+ionomycin stimulated C57BL/6 mouse splenocytes (6 hours) stained with anti-CD3 PE (17A2) and intracellularly stained with JES6-5H4 APC

## Applications:

**Applications:** ICFC - Quality tested

**Recommended Usage:** Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.25 µg per 10<sup>6</sup> cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** **ELISA Detection<sup>1-3</sup> or ELISPOT Detection<sup>4-6</sup>:** The biotinylated JES6-5H4 antibody is useful as a detection antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with the purified JES6-1A12 antibody (Cat. No. 503702/503704) as capture antibody and recombinant mouse IL-2 (Cat. No. 575409) as the standard. **Flow Cytometry<sup>8-10</sup>:** The fluorochrome-labeled JES6-5H4 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IL-2-producing cells within mixed cell populations. **Neutralization<sup>1,7</sup>:** The LEAF™ purified antibody (Endotoxin in vivo and *in vitro* (Cat. No. 503812) is recommended for neutralization.

**Additional reported applications (for the relevant formats) include:** immunoprecipitation<sup>1</sup>, immunohistochemical staining of paraformaldehyde-fixed, saponin-treated frozen tissue sections<sup>2</sup>, *in vivo* capture<sup>7</sup>, and immunocytochemistry.

**Note:** For testing mouse IL-2 in serum, plasma or supernatant, BioLegend's ELISA Max™ Sets (Cat. No. 431001 to 431006) are specially developed and recommended.

- Application References:**
- Abrams J, *et al.* 1992. *Immunol. Rev.* 127:5.
  - Sander B, *et al.* 1993. *J. Immunol. Meth.* 166:201.
  - Abrams J. 1995. *Curr. Prot. Immunol.* John Wiley and Sons New York. Unit 6.20.
  - Klinman D, *et al.* 1994. *Curr. Prot. Immunol.* John Wiley and Sons New York. Unit 6.19.
  - Mo X, *et al.* 1995. *J. Virol.* 69:1288.
  - Karulin A, *et al.* 2000. *J. Immunol.* 164:1862.
  - Finkelman F, *et al.* 2003. *Curr. Prot. Immunol.* John Wiley & Sons New York. Unit 6.28.
  - Ko SY, *et al.* 2005. *J. Immunol.* 175:3309. PubMed
  - Kang SS and Allen PM. 2005. *J. Immunol.* 174:5382.
  - Lawson BR, *et al.* 2007. *J. Immunol.* 178:5366.
  - Lei F, *et al.* 2012. *J Vis exp.* 60:3986. PubMed.

**Description:** IL-2 is a potent lymphoid cell growth factor which exerts its biological activity primarily on T cells. Additionally, IL-2 has been found to stimulate growth and differentiation of B cells, NK cells, LAK cells, monocytes, and oligodendrocytes.

- Antigen References:**
- Fitzgerald K, *et al.* Eds. 2001. The Cytokine FactsBook. Academic Press San Diego.
  - Taniguchi T, *et al.* 1993. *Cell* 73:5.
  - Nisticò G. 1993. *Prog. Neurobiol.* 40:463.
  - Waldmann T, *et al.* 1993. *Ann. NY Acad. Sci.* 685:603.

Related Products:	Product	Clone	Application
	Cell Staining Buffer		FC, ICC, ICFC
	Fixation Buffer		ICC, ICFC
	Permeabilization Wash Buffer (10X)		ICC, ICFC, IHC
	Brefeldin A Solution (1,000X)		ICFC
	Monensin Solution (1,000X)		ICFC
	RBC Lysis Buffer (10X)		FC, ICFC
	APC Rat IgG2b, κ Isotype Ctrl	RTK4530	FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.