

Product Data Sheet

LEAF™ Purified anti-human IL-12/IL-23 p40 (monomer, dimer, heterodimer)

Catalog # / Size: 501813 / 500 µg

Clone: C11.5

Isotype: Mouse IgG1, κ

Immunogen: CHO-expressed, recombinant human IL-12 p70 heterod

Reactivity: Human, IL-12/IL-23 p40 subunit (monomer, homodimer and heterodimer IL-12 p35/p40 or IL-23 p19/p40)

Preparation: The LEAF™ (Low Endotoxin, Azide-Free) antibody was purified by affinity chromatography.

Formulation: 0.2 µm filtered in phosphate-buffered solution, pH 7.2, containing no preservative. Endotoxin level is <0.1 EU/µg of

the protein (<0.01 ng/µg of the protein) as determined by the LAL test.

Concentration: 1.0 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C. This LEAF™ solution contains no preservative; handle under

aseptic conditions.

Applications:

Applications: ICFC - Quality tested

Block, IHC, WB, Neut - Reported in the literature

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric

analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.125 µg per million cells in 100 µl volume or 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each

application.

Application Notes: Flow Cytometry^{3,5}: The fluorochrome-labeled C11.5 antibody is useful for intracellular immunofluorescent staining

and flow cytometric analysis to identify IL-12 p40 (monomeric, heterodimeric, homodimeric) -producing cells within mixed cell populations. For intracellular cytokine staining protocol, please visit www.biolegend.com and click on the

Neutralization⁶: The LEAF™ purified antibody (Endotoxin Additional reported applications (for the relevant formats) include: Western blotting, ELISA Capture^{1,2}, ICFC Blocking, immunohistochemical staining⁴ of

paraformaldehyde-fixed, saponin-treated tissue frozen sections, and immunocytochemistry.

Note: For testing human IL-12/IL-23 p40 (monomer, dimer) in serum or plasma, BioLegend's ELISA Max™ Sets (Cat. No. 430701 to 430706) are specially developed and recommended. For testing human IL-12 p70 in serum or plasma, BioLegend's ELISA Max™ Sets (Cat. No. 431701 to 431706) are specially developed and recommended.

Application References: 1. D'Andrea A, et al. 1992. J. Exp. Med. 176:1387. 2. D'Andrea A, et al. 1993. J. Exp. Med. 178:1041. 3. Prussin C, et al. 1995. J. Immunol. Meth. 188:117.

4. Leifeld L, et al. 2002. Hepatology 36:1001.

5. Pekkari K, et al. 2001. Blood 97:3184.

6. Bolovan-Fritts CA, et al. 2004. J. Virol. 78:13173.

Description: The C11.5 antibody reacts with human IL-12 p40 subunit of the IL-12 p70 heterodimer and IL-23 p40 subunit of the

IL-23 p19/p40, as well as p40 monomer and homodimer. The C11.5 antibody can neutralize the bioactivity of natural

or recombinant IL-12 p70.

Antigen References: 1. Fitzgerald K, et al. Eds. 2001. The Cytokine FactsBook. Academic Press San Diego.

Quesniaux V. 1992. Research Immunol. 143:385.
Trinchieri G, et al. 1992 Prog. Growth Factor Res. 4:355.

4. Trinchieri G, et al. 1993 Immunol. Today. 14:335.

Oppmann B, *et al.* 2000 *Immunity*. 13:715.
Aggarwal S, *et al.* 2003 *J. Biol. Chem.* 278:1910.

7. Parham C, et al. 2002 J. Immunol. 168:5699.

8. Belladonna ML, et al. 2002 J. Immunol. 168:5448. 9. Lankford CS, et al. 2003 J. Leukoc. Biol. 73:49.

Related Products: Product Clone Application

LEAF™ Purified Mouse IgG1, κ Isotype Ctrl

Recombinant Human IL-12/IL-23 (p40) Recombinant Human IL-12 (p70)

FC, ICFC, WB, IP, ICC, IF, FA MOPC-21 **ELISA**

BA, ELISA



