

Purified anti-human IL-13

Catalog # / Size: 501901 / 50 µg
501902 / 500 µg

Clone: JES10-5A2

Isotype: Rat IgG1, κ

Immunogen: COS-expressed, recombinant human IL-13


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Applications:

Applications: ELISA Capture - *Quality tested*
IHC, WB - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by ELISA analysis. For ELISA applications, a concentration range of 1-4 µg/ml is recommended. To obtain a linear standard curve, serial dilutions of human IL-13 recombinant protein ranging from 2000 to 8 pg/ml are recommended for each ELISA plate. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: **ELISA or ELISPOT Capture^{1,2}:** The purified JES10-5A2 antibody is useful as the capture antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the biotinylated Poly5020 antibody (Cat. No. 502001) as the detecting antibody. The LEAF™ purified antibody is suggested for ELISPOT capture.

Application References:

1. McKenzie, A., *et al.* 1994. *Curr. Prot. Immunol.* John Wiley & Sons. New York. Section 6.18.
2. Abrams, J. 1995. *Curr. Prot. Immunol.* John Wiley and Sons, New York. Unit 6.20.
3. Schaerli, P., *et al.* 2000. *J. Exp. Med.* 192:1553.
4. Andersson, J., *et al.* 1994. *Immunology* 83:16.
5. Skansen-Saphir, U., *et al.* 1994. *Eur. J. Immunol.* 24:916.
6. Andersson, U., *et al.* 1999. *Detection and quantification of gene expression.* New York:Springer-Verlag.
7. Gelfand, E. W., *et al.* 2006. *J. Aller. and Clinic. Immunol.* 117:577.PubMed
8. Knorr, C., *et al.* 1999. *Am. J. Pathol.* 155:2019.

Description: IL-13 is an immunoregulatory cytokine produced primarily by activated Th2 lymphocytes. IL-13 shares 30% amino acid sequence homology with IL-4 and demonstrates similar biological activities. The biological activities of IL-13 include: suppression of macrophage cytotoxic activity, upregulation of IL-1RA expression, and suppression of proinflammatory cytokine secretion. The JES10-5A2 antibody reacts with human interleukin-13 (IL-13). The JES10-5A2 antibody can neutralize the bioactivity of natural or recombinant IL-13.

Antigen References:

1. Fitzgerald, K., *et al.* Eds. 2001. *The Cytokine FactsBook.* Academic Press, San Diego.
2. Hilton, D., *et al.* 1996. *P. Natl. Acad. Sci. USA* 93:497.
3. Obiri, N., *et al.* 1995. *J. Biol. Chem.* 270:8797.
4. Ochensberger, B., *et al.* 1996. *Blood* 88:3028.

Related Products:	Product	Clone	Application
	Biotin anti-human IL-13	Poly5020	ELISA Detection, ELISPOT
	Recombinant Human IL-13	rh IL-13	Detection
	HRP Avidin	Avidin	BA, ELISA
	TMB Substrate Reagent Set		ELISA, ELISPOT, IHC, WB
	ELISA Assay Diluent (5X)		ELISA
			ELISA



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.